

# *Sharp Trace Hardy–Sobolev–Maz’ya Inequalities and the Fractional Laplacian*

**Stathis Filippas, Luisa Moschini &  
Achilles Tertikas**

**Archive for Rational Mechanics and  
Analysis**

ISSN 0003-9527


Volume 208

Number 1

Arch Rational Mech Anal (2013)

208:109-161

DOI 10.1007/s00205-012-0594-4



**Your article is protected by copyright and all rights are held exclusively by Springer-Verlag Berlin Heidelberg. This e-offprint is for personal use only and shall not be self-archived in electronic repositories. If you wish to self-archive your work, please use the accepted author's version for posting to your own website or your institution's repository. You may further deposit the accepted author's version on a funder's repository at a funder's request, provided it is not made publicly available until 12 months after publication.**

# Sharp Trace Hardy–Sobolev–Maz’ya Inequalities and the Fractional Laplacian

STATHIS FILIPPAS, LUISA MOSCHINI & ACHILLES TERTIKAS

Communicated by F. LIN

## Abstract

In this work we establish trace Hardy and trace Hardy–Sobolev–Maz’ya inequalities with best Hardy constants for domains satisfying suitable geometric assumptions such as mean convexity or convexity. We then use them to produce fractional Hardy–Sobolev–Maz’ya inequalities with best Hardy constants for various fractional Laplacians. In the case where the domain is the half space, our results cover the full range of the exponent  $s \in (0, 1)$  of the fractional Laplacians. In particular, we give a complete answer in the  $L^2$  setting to an open problem raised by FRANK and SEIRINGER (“Sharp fractional Hardy inequalities in half-spaces,” in *Around the research of Vladimir Maz’ya*. International Mathematical Series, 2010).

## 1. Introduction

The Hardy inequality in the upper half space asserts that

$$\int_{\mathbf{R}_+^n} |\nabla u|^2 \, dx \geq \frac{1}{4} \int_{\mathbf{R}_+^n} \frac{|u|^2}{x_n^2} \, dx, \quad u \in C_0^\infty(\mathbf{R}_+^n), \quad (1.1)$$

where  $\mathbf{R}_+^n = \{(x_1, \dots, x_n) : x_n > 0\}$  denotes the upper half-space and  $\frac{1}{4}$  is the best possible constant.

If  $\Omega \subset \mathbf{R}^n$  and  $d(x) = \text{dist}(x, \partial\Omega)$ , then there are two main directions towards establishing Hardy inequalities. One direction is to find proper regularity assumptions on the boundary of  $\Omega$  that imply the existence of a positive constant  $C_\Omega$ , such that

$$\int_{\Omega} |\nabla u|^2 \, dx \geq C_\Omega \int_{\Omega} \frac{|u|^2}{d^2(x)} \, dx, \quad u \in C_0^\infty(\Omega).$$

In this direction we refer to [4, 29] and references therein.

A second direction aims at finding geometric assumptions on  $\Omega$  that imply the Hardy inequality with best constant  $\frac{1}{4}$ , that is

$$\int_{\Omega} |\nabla u|^2 dx \geq \frac{1}{4} \int_{\Omega} \frac{|u|^2}{d^2(x)} dx, \quad u \in C_0^\infty(\Omega). \tag{1.2}$$

The standard geometric assumption here is the convexity of  $\Omega$ , see, for example, [10, 18, 19]. However, inequality (1.2) remains true under the weaker assumption

$$-\Delta d(x) \geq 0, \quad x \in \Omega. \tag{1.3}$$

This is meant in the distributional sense. We refer to [6] where this condition arises in a natural way. In fact, condition (1.3) is equivalent to convexity in two space dimensions, but it is weaker than convexity for  $n \geq 3$ , since any convex domain satisfies (1.3), whereas there are nonconvex domains that satisfy (1.3) [5]. We emphasize that there is no need for further regularity assumptions on  $\Omega$ . In the case where  $\partial\Omega$  is  $C^2$ , condition (1.3) has recently been shown to be equivalent to the mean convexity of  $\partial\Omega$ , that is,  $(n - 1)H(x) = -\Delta d(x) \geq 0$  for  $x \in \partial\Omega$ , see [31, 37].

If in addition to (1.3) the domain  $\Omega$  is a  $C^2$  domain with finite inner radius, then it has been established that one can combine the Sobolev and the Hardy inequalities, the latter with best constant. More precisely, for  $n \geq 3$  there exists a positive constant  $c$  such that

$$\int_{\Omega} |\nabla u|^2 dx \geq \frac{1}{4} \int_{\Omega} \frac{|u|^2}{d^2(x)} dx + c \left( \int_{\Omega} |u|^{\frac{2n}{n-2}} dx \right)^{\frac{n-2}{n}}, \quad u \in C_0^\infty(\Omega), \tag{1.4}$$

see [23]. In [27] Hardy–Sobolev–Maz’ya inequalities are established under a different geometric assumption than (1.3), which allows infinite inner radius. Frank and Loss established in [26] inequality (1.4) with a constant  $c$  independent of  $\Omega$ , when  $\Omega$  is convex.

Recently, much attention has been devoted to the fractional Laplacian, which for  $s \in (0, 1)$  is defined as

$$(-\Delta)^s f(x) = c_{n,s} \text{PV} \int_{\mathbf{R}^n} \frac{f(x) - f(\xi)}{|x - \xi|^{n+2s}} d\xi, \tag{1.5}$$

where PV stands for the Cauchy principal value and

$$c_{n,s} = \frac{s 2^{2s} \Gamma\left(\frac{n+2s}{2}\right)}{\Gamma(1-s)\pi^{\frac{n}{2}}}. \tag{1.6}$$

There are other ways to define the fractional Laplacian, as, for instance, via the Fourier transform. We note that the fractional Laplacian is a nonlocal operator and that this raises several technical difficulties. However, there is a way of studying various properties of the fractional Laplacian via the Dirichlet to Neumann map. This has been recently studied by CAFFARELLI and SILVESTRE [13], and it will be central in this work. Let us briefly recall the approach in [13] where, by adding

a new variable  $y$ , they relate the fractional Laplacian to a local operator. For any function  $f$  one solves the following extension problem

$$\operatorname{div}(y^{1-2s}\nabla_{(x,y)}u(x,y)) = 0, \quad \mathbf{R}^n \times (0, \infty), \tag{1.7}$$

$$u(x, 0) = f(x), \quad \mathbf{R}^n, \tag{1.8}$$

the natural energy of which is given by

$$J[u] = \int_0^{+\infty} \int_{\mathbf{R}^n} y^{1-2s} |\nabla_{(x,y)}u(x,y)|^2 \, dx \, dy.$$

Then, up to a normalizing factor  $C$  one establishes that

$$-\lim_{y \rightarrow 0^+} y^{1-2s} u_y(x,y) = C(-\Delta)^s f(x).$$

Our interest in this work is to study the fractional Laplacian defined in subsets of  $\mathbf{R}^n$  and, in particular, to establish Hardy and Hardy–Sobolev–Maz’ya inequalities there. There is much interest in fractional Laplacian in subsets of  $\mathbf{R}^n$  coming from various applications, as, for instance, censored stable processes and killed stable processes [7, 15–17], Gamma convergence and phase transition problems [2, 28, 36, 38, 39] and nonlinear PDE theory [11, 12, 42]. In [8] it was conjectured that the best Hardy constant in the case of the fractional Laplacian associated to a censored stable process is the same for all convex domains. In [24] the question was posed regarding establishing fractional Hardy–Sobolev–Maz’ya inequalities for the half space.

Contrary to the case of the full space  $\mathbf{R}^n$ , there are several different fractional Laplacians that one can define on a domain  $\Omega \subset \mathbf{R}^n$ . In particular, in the above mentioned references three different fractional Laplacians appear. In all cases we will use the Dirichlet to Neumann map after identifying the proper extension problem. Throughout this work we assume that the domain  $\Omega$  is a uniformly Lipschitz domain; for the precise definition see Section 2.

We start with the fractional Laplacian that appears in [11, 12, 42]. The proper extension problem in this case is to consider test functions in  $C_0^\infty(\Omega \times \mathbf{R})$ . At this point we recall that the inner radius of a domain  $\Omega$  is defined as  $R_{\text{in}} := \sup_{x \in \Omega} d(x)$ . We say that the domain  $\Omega$  has a finite inner radius whenever  $R_{\text{in}} < \infty$ . Our first result concerns the extended problem and reads:

**Theorem 1.** (trace Hardy and trace Hardy–Sobolev–Maz’ya I) *Let  $\frac{1}{2} \leq s < 1$ ,  $n \geq 2$  and  $\Omega \subset \mathbf{R}^n$  be a domain.*

(i) *If, in addition,  $\Omega$  is such that*

$$-\Delta d(x) \geq 0, \quad x \in \Omega, \tag{1.9}$$

*then for all  $u \in C_0^\infty(\Omega \times \mathbf{R})$  there holds*

$$\int_0^{+\infty} \int_{\Omega} y^{1-2s} |\nabla_{(x,y)}u(x,y)|^2 \, dx \, dy \geq \bar{d}_s \int_{\Omega} \frac{u^2(x,0)}{d^{2s}(x)} \, dx, \tag{1.10}$$

with

$$\bar{d}_s := \frac{2\Gamma(1-s)\Gamma^2\left(\frac{3+2s}{4}\right)}{\Gamma^2\left(\frac{3-2s}{4}\right)\Gamma(s)}. \tag{1.11}$$

(ii) Suppose there exists a point  $x_0 \in \partial\Omega$  and  $r > 0$  such that the part of the boundary  $\partial\Omega \cap B(x_0, r)$  is  $C^1$  regular. Then

$$\bar{d}_s \geq \inf_{u \in C_0^\infty(\Omega \times \mathbf{R})} \frac{\int_0^{+\infty} \int_\Omega y^{1-2s} |\nabla u|^2 \, dx \, dy}{\int_\Omega \frac{u^2(x,0)}{d^{2s}(x)} \, dx}.$$

In particular,  $\bar{d}_s$  in (1.10) is the best constant.

(iii) If  $\Omega$  is a uniformly Lipschitz domain with finite inner radius satisfying (1.9), and  $s \in (\frac{1}{2}, 1)$ , then there exists a positive constant  $c$  such that for  $u \in C_0^\infty(\Omega \times \mathbf{R})$  there holds

$$\begin{aligned} \int_0^{+\infty} \int_\Omega y^{1-2s} |\nabla_{(x,y)} u(x, y)|^2 \, dx \, dy &\geq \bar{d}_s \int_\Omega \frac{u^2(x, 0)}{d^{2s}(x)} \, dx \\ &\quad + c \left( \int_\Omega |u(x, 0)|^{\frac{2n}{n-2s}} \, dx \right)^{\frac{n-2s}{n}}. \end{aligned} \tag{1.12}$$

Actually, in the case of half space  $\Omega = \mathbf{R}_+^n$  we establish a much stronger result covering the full range  $s \in (0, 1)$ . In particular, we have

**Theorem 2.** (half space, trace Hardy–Sobolev–Maz’ya I) *Let  $0 < s < 1$  and  $n \geq 2$ .*

(i) *For all  $u \in C_0^\infty(\mathbf{R}_+^n \times \mathbf{R})$  there holds*

$$\int_0^\infty \int_{\mathbf{R}_+^n} y^{1-2s} |\nabla_{(x,y)} u(x, y)|^2 \, dx \, dy \geq \bar{d}_s \int_{\mathbf{R}_+^n} \frac{u^2(x, 0)}{x_n^{2s}} \, dx, \tag{1.13}$$

with

$$\bar{d}_s := \frac{2\Gamma(1-s)\Gamma^2\left(\frac{3+2s}{4}\right)}{\Gamma^2\left(\frac{3-2s}{4}\right)\Gamma(s)}. \tag{1.14}$$

(ii) *The constant  $\bar{d}_s$  in (1.13) is sharp, that is,*

$$\bar{d}_s = \inf_{u \in C_0^\infty(\mathbf{R}_+^n \times \mathbf{R})} \frac{\int_0^\infty \int_{\mathbf{R}_+^n} y^{1-2s} |\nabla u|^2 \, dx \, dy}{\int_{\mathbf{R}_+^n} \frac{u^2(x,0)}{x_n^{2s}} \, dx}.$$

(iii) *There exists a positive constant  $c$  such that for all  $u \in C_0^\infty(\mathbf{R}_+^n \times \mathbf{R})$  there holds*

$$\begin{aligned} \int_0^\infty \int_{\mathbf{R}_+^n} y^{1-2s} |\nabla_{(x,y)} u(x, y)|^2 \, dx \, dy &\geq \bar{d}_s \int_{\mathbf{R}_+^n} \frac{u^2(x, 0)}{x_n^{2s}} \, dx \\ &\quad + c \left( \int_{\mathbf{R}_+^n} |u(x, 0)|^{\frac{2n}{n-2s}} \, dx \right)^{\frac{n-2s}{n}}. \end{aligned} \tag{1.15}$$

We will apply Theorem 1 to the fractional Laplacian that is defined as follows. Let  $\Omega \subset \mathbf{R}^n$  be a bounded domain, and let  $\lambda_i$  and  $\phi_i$  be the Dirichlet eigenvalues and orthonormal eigenfunctions of the Laplacian, that is,  $-\Delta\phi_i = \lambda_i\phi_i$  in  $\Omega$ , with  $\phi_i = 0$  on  $\partial\Omega$ . Then, for  $f(x) = \sum c_i\phi_i(x)$  we define

$$(-\Delta)^s f(x) = \sum_{i=1}^{\infty} c_i \lambda_i^s \phi_i(x), \quad 0 < s < 1, \tag{1.16}$$

in which case

$$((-\Delta)^s f, f)_{\Omega} = \int_{\Omega} f(x)(-\Delta)^s f(x) \, dx = \sum_{i=1}^{\infty} c_i^2 \lambda_i^s. \tag{1.17}$$

In the sequel we will refer to this fractional Laplacian as the *spectral fractional Laplacian*. We then have

**Theorem 3.** (Hardy and Hardy–Sobolev–Maz’ya for spectral fractional Laplacian) *Let  $\frac{1}{2} \leq s < 1$ ,  $n \geq 2$  and  $\Omega \subset \mathbf{R}^n$  be a bounded domain.*

(i) *If, in addition,  $\Omega$  is such that*

$$-\Delta d(x) \geq 0, \quad x \in \Omega, \tag{1.18}$$

*then, for all  $f \in C_0^\infty(\Omega)$  there holds*

$$((-\Delta)^s f, f)_{\Omega} \geq d_s \int_{\Omega} \frac{f^2(x)}{d^{2s}(x)} \, dx, \tag{1.19}$$

*with*

$$d_s := \frac{2^{2s} \Gamma^2\left(\frac{3+2s}{4}\right)}{\Gamma^2\left(\frac{3-2s}{4}\right)}. \tag{1.20}$$

(ii) *Suppose there exists a point  $x_0 \in \partial\Omega$  and  $r > 0$  such that the part of the boundary  $\partial\Omega \cap B(x_0, r)$  is  $C^1$  regular. Then*

$$d_s \geq \inf_{f \in C_0^\infty(\Omega)} \frac{((-\Delta)^s f, f)_{\Omega}}{\int_{\Omega} \frac{f^2(x)}{d^{2s}(x)} \, dx}.$$

(iii) *If  $\Omega$  is a Lipschitz domain satisfying (1.18) and  $s \in (\frac{1}{2}, 1)$ , then there exists a positive constant  $c$  such that for all  $f \in C_0^\infty(\Omega)$  there holds*

$$((-\Delta)^s f, f)_{\Omega} \geq d_s \int_{\Omega} \frac{f^2(x)}{d^{2s}(x)} \, dx + c \left( \int_{\Omega} |f(x)|^{\frac{2n}{n-2s}} \, dx \right)^{\frac{n-2s}{n}}. \tag{1.21}$$

We next consider the fractional Laplacian associated to the killed stable processes that appears in [7,8,36,38,39], which from now on we will call the *Dirichlet fractional Laplacian*. The proper extension problem involves test functions  $u \in C_0^\infty(\mathbf{R}^n \times \mathbf{R})$  such that  $u(x, 0) = 0$  in the complement of  $\Omega$ , that is, for  $x \in \mathcal{C}\Omega$ . For this fractional Laplacian, our assumption on the domain  $\Omega$  is convexity instead of (1.3). The reason for this is that our method requires subharmonicity of the distance function in  $\mathcal{C}\Omega$ , which is equivalent to the convexity of  $\Omega$ , see [5]. Our next result reads:

**Theorem 4.** (trace Hardy and trace Hardy–Sobolev–Maz’ya II) *Let  $\frac{1}{2} \leq s < 1$ ,  $n \geq 2$  and  $\Omega \subset \mathbf{R}^n$  be a domain.*

(i) *If, in addition,  $\Omega$  is convex, then for all  $u \in C_0^\infty(\mathbf{R}^n \times \mathbf{R})$  such that  $u(x, 0) = 0$  for  $x \in \mathcal{C}\Omega$ , there holds*

$$\int_0^{+\infty} \int_{\mathbf{R}^n} y^{1-2s} |\nabla_{(x,y)} u(x, y)|^2 dx dy \geq \bar{k}_s \int_{\Omega} \frac{u^2(x, 0)}{d^{2s}(x)} dx, \quad (1.22)$$

with

$$\bar{k}_s := \frac{2^{1-2s} \Gamma^2(s + \frac{1}{2}) \Gamma(1 - s)}{\pi \Gamma(s)}. \quad (1.23)$$

(ii) *Suppose there exists a point  $x_0 \in \partial\Omega$  and  $r > 0$  such that the part of the boundary  $\partial\Omega \cap B(x_0, r)$  is  $C^1$  regular. Then*

$$\bar{k}_s \geq \inf_{\substack{u \in C_0^\infty(\mathbf{R}^n \times \mathbf{R}), \\ u(x, 0) = 0, x \in \mathcal{C}\Omega}} \frac{\int_0^{+\infty} \int_{\mathbf{R}^n} y^{1-2s} |\nabla u|^2 dx dy}{\int_{\Omega} \frac{u^2(x, 0)}{d^{2s}(x)} dx}.$$

In particular,  $\bar{k}_s$  in (1.22) is the best constant.

(iii) *If  $\Omega$  is a uniformly Lipschitz and convex domain with finite inner radius and  $s \in (\frac{1}{2}, 1)$ , then there exists a positive constant,  $c$ , such that the following improvement holds true for all  $u \in C_0^\infty(\mathbf{R}^n \times \mathbf{R})$  with  $u(x, 0) = 0$  for  $x \in \mathcal{C}\Omega$ :*

$$\begin{aligned} \int_0^{+\infty} \int_{\mathbf{R}^n} y^{1-2s} |\nabla_{(x,y)} u(x, y)|^2 dx dy &\geq \bar{k}_s \int_{\Omega} \frac{u^2(x, 0)}{d^{2s}(x)} dx \\ &+ c \left( \int_{\Omega} |u(x, 0)|^{\frac{2n}{n-2s}} dx \right)^{\frac{n-2s}{n}}. \end{aligned} \quad (1.24)$$

Elementary manipulations show that

$$\bar{d}_s = 2 \sin^2 \left( \frac{(2s + 1)\pi}{4} \right) \bar{k}_s,$$

thus

$$\bar{d}_s > \bar{k}_s, \quad \text{for } s \in (0, 1),$$

which implies, in particular, that the best constants of Theorems 1 and 4 are different.

We next apply Theorem 4 to the Dirichlet fractional Laplacian. In this case, for  $f \in C_0^\infty(\Omega)$  we extend  $f$  in all of  $\mathbf{R}^n$  by setting  $f = 0$  in  $\mathcal{C}\Omega$  and use (1.5). In particular, the corresponding quadratic form is

$$\begin{aligned} ((-\Delta)_D^s f, f)_{\mathbf{R}^n} &= \frac{c_{n,s}}{2} \int_{\mathbf{R}^n} \int_{\mathbf{R}^n} \frac{|f(x) - f(\xi)|^2}{|x - \xi|^{n+2s}} dx d\xi \\ &= \frac{c_{n,s}}{2} \left( \int_{\Omega} \int_{\Omega} \frac{|f(x) - f(\xi)|^2}{|x - \xi|^{n+2s}} dx d\xi + 2 \int_{\Omega} \int_{\mathcal{C}\Omega} \frac{|f(x)|^2}{|x - \xi|^{n+2s}} dx d\xi \right), \end{aligned} \quad (1.25)$$

with the constant  $c_{n,s}$  as given by Equation (1.6). We then have:



**Theorem 5.** (Hardy and Hardy–Sobolev–Maz’ya for Dirichlet fractional Laplacian) *Let  $\frac{1}{2} \leq s < 1$ ,  $n \geq 2$  and  $\Omega \subset \mathbf{R}^n$  be a domain.*

(i) *If, in addition,  $\Omega$  is convex, then for all  $f \in C_0^\infty(\Omega)$  there holds*

$$((-\Delta)_D^s f, f)_{\mathbf{R}^n} \geq \frac{\Gamma^2(s + \frac{1}{2})}{\pi} \int_{\Omega} \frac{f^2(x)}{d^{2s}(x)} dx. \tag{1.26}$$

*Equivalently, one has that*

$$\int_{\mathbf{R}^n} \int_{\mathbf{R}^n} \frac{|f(x) - f(\xi)|^2}{|x - \xi|^{n+2s}} dx d\xi \geq k_{n,s} \int_{\Omega} \frac{f^2(x)}{d^{2s}(x)} dx, \tag{1.27}$$

where

$$k_{n,s} := \frac{2^{1-2s} \pi^{\frac{n-2}{2}} \Gamma(1-s) \Gamma^2(s + \frac{1}{2})}{s \Gamma(\frac{n+2s}{2})}. \tag{1.28}$$

(ii) *Suppose there exists a point  $x_0 \in \partial\Omega$  and  $r > 0$  such that the part of the boundary  $\partial\Omega \cap B(x_0, r)$  is  $C^1$  regular. Then the Hardy constants  $\frac{\Gamma^2(s + \frac{1}{2})}{\pi}$  in (1.26) and  $k_{n,s}$  in (1.27) are optimal.*

(iii) *If  $\Omega$  is a uniformly Lipschitz and convex domain with finite inner radius and  $s \in (\frac{1}{2}, 1)$ , then there exists a positive constant  $c$  such that for all  $f \in C_0^\infty(\Omega)$  there holds*

$$\begin{aligned} ((-\Delta)_D^s f, f)_{\mathbf{R}^n} &\geq \frac{\Gamma^2(s + \frac{1}{2})}{\pi} \int_{\Omega} \frac{f^2(x)}{d^{2s}(x)} dx \\ &\quad + c \left( \int_{\Omega} |f(x)|^{\frac{2n}{n-2s}} dx \right)^{\frac{n-2s}{n}}. \end{aligned} \tag{1.29}$$

*Equivalently, one has that*

$$\begin{aligned} \int_{\mathbf{R}^n} \int_{\mathbf{R}^n} \frac{|f(x) - f(\xi)|^2}{|x - \xi|^{n+2s}} dx d\xi &\geq k_{n,s} \int_{\Omega} \frac{f^2(x)}{d^{2s}(x)} dx \\ &\quad + c \left( \int_{\Omega} |f(x)|^{\frac{2n}{n-2s}} dx \right)^{\frac{n-2s}{n}}. \end{aligned} \tag{1.30}$$

The case where  $\Omega$  is the half-space  $\Omega = \mathbf{R}_+^n = \{(x_1, \dots, x_n) : x_n > 0\}$  is of particular interest, see [7, 8, 21, 24, 40]. In this case we obtain a stronger result that covers the full range  $s \in (0, 1)$ . More precisely, we have:

**Theorem 6.** (half space, trace Hardy–Sobolev–Maz’ya and fractional Hardy–Sobolev–Maz’ya II) *Let  $0 < s < 1$  and  $n \geq 2$ .*

(i) *For all  $u \in C_0^\infty(\mathbf{R}^n \times \mathbf{R})$  with  $u(x, 0) = 0$ ,  $x \in \mathbf{R}_+^n$ , there holds*

$$\int_0^{+\infty} \int_{\mathbf{R}_+^n} y^{1-2s} |\nabla_{(x,y)} u(x, y)|^2 dx dy \geq \bar{k}_s \int_{\mathbf{R}_+^n} \frac{u^2(x, 0)}{x_n^{2s}} dx, \tag{1.31}$$

where

$$\bar{k}_s := \frac{2^{1-2s} \Gamma^2(s + \frac{1}{2}) \Gamma(1 - s)}{\pi \Gamma(s)},$$

is the best constant in (1.31).

(ii) There exists a positive constant  $c$ , such that for all  $u \in C_0^\infty(\mathbf{R}^n \times \mathbf{R})$  with  $u(x, 0) = 0, x \in \mathbf{R}_-^n$ , there holds

$$\begin{aligned} \int_0^{+\infty} \int_{\mathbf{R}^n} y^{1-2s} |\nabla_{(x,y)} u(x, y)|^2 dx dy &\geq \bar{k}_s \int_{\mathbf{R}_+^n} \frac{u^2(x, 0)}{x_n^{2s}} dx \\ &+ c \left( \int_{\mathbf{R}_+^n} |u(x, 0)|^{\frac{2n}{n-2s}} dx \right)^{\frac{n-2s}{n}}. \end{aligned} \tag{1.32}$$

(iii) As a consequence, there exists a positive constant  $c$  such that for all  $f \in C_0^\infty(\mathbf{R}_+^n)$  there holds

$$\begin{aligned} \int_{\mathbf{R}^n} \int_{\mathbf{R}^n} \frac{|f(x) - f(\xi)|^2}{|x - \xi|^{n+2s}} dx d\xi &\geq k_{n,s} \int_{\mathbf{R}_+^n} \frac{f^2(x)}{x_n^{2s}} dx \\ &+ c \left( \int_{\mathbf{R}_+^n} |f(x)|^{\frac{2n}{n-2s}} dx \right)^{\frac{n-2s}{n}}, \end{aligned} \tag{1.33}$$

where  $k_{n,s}$  is given by Equation (1.28).

Or, equivalently, for all  $f \in C_0^\infty(\mathbf{R}_+^n)$  there holds

$$\begin{aligned} \int_{\mathbf{R}_+^n} \int_{\mathbf{R}_+^n} \frac{|f(x) - f(\xi)|^2}{|x - \xi|^{n+2s}} dx d\xi &\geq \kappa_{n,s} \int_{\mathbf{R}_+^n} \frac{f^2(x)}{x_n^{2s}} dx \\ &+ c \left( \int_{\mathbf{R}_+^n} |f(x)|^{\frac{2n}{n-2s}} dx \right)^{\frac{n-2s}{n}}, \end{aligned} \tag{1.34}$$

where

$$\kappa_{n,s} := \pi^{\frac{n-1}{2}} \frac{\Gamma(s + \frac{1}{2})}{s \Gamma(\frac{n+2s}{2})} \left[ \frac{2^{1-2s}}{\sqrt{\pi}} \Gamma(1 - s) \Gamma\left(s + \frac{1}{2}\right) - 1 \right].$$

We note that the Hardy–Sobolev–Maz’ya inequality (1.33) refers to the Dirichlet fractional Laplacian associated to the killed stable processes, whereas inequality (1.34) is associated to the censored stable processes. The Hardy constants  $k_{n,s}$  and  $\kappa_{n,s}$  appearing in (1.33) and (1.34), respectively, are optimal, as shown in [8]. The corresponding fractional Hardy inequality of (1.34) with best constant, in the case of a convex domain  $\Omega$ , that is,

$$\int_{\Omega} \int_{\Omega} \frac{|f(x) - f(\xi)|^2}{|x - \xi|^{n+2s}} dx d\xi \geq \kappa_{n,s} \int_{\Omega} \frac{f^2(x)}{d(x)^{2s}} dx, \quad f \in C_0^\infty(\Omega),$$

has been established for  $s \in (\frac{1}{2}, 1)$  in [32]. The question of obtaining a Hardy–Sobolev–Maz’ya inequality for the half space was raised in [24] and was answered positively in the  $L^2$  setting in [21, 40], but only for the range  $s \in (\frac{1}{2}, 1)$ .

For other types of trace Hardy inequalities we refer to [3, 20]. We finally note that fractional Sobolev inequalities play an important role in many other directions, see for example, [9, 14, 34, 35].

## 2. The Trace Hardy Inequality I

In this section we will prove the trace Hardy inequality contained in Theorem 1. We first recall the definition of a uniformly Lipschitz domain  $\Omega$ ; see section 12 of [30]. We note that Stein calls such a domain minimally smooth, see section 3.3 of [41].

A domain  $\Omega$  is called uniformly Lipschitz if there exist  $\varepsilon > 0$ ,  $L > 0$ , and  $M \in \mathbf{N}$  and a locally finite countable cover  $\{U_i\}$  of  $\partial\Omega$  with the following properties:

- (i) If  $x \in \partial\Omega$  then  $B(x, \varepsilon) \subset U_i$  for some  $i$ .
- (ii) Every point of  $\mathbf{R}^n$  is contained in at most  $M$   $U_i$ 's.
- (iii) For each  $i$  there exist local coordinates  $y = (y', y_n) \in \mathbf{R}^{n-1} \times \mathbf{R}$  and a Lipschitz function  $f : \mathbf{R}^{n-1} \rightarrow \mathbf{R}$ , with  $\text{Lip}f \leq L$  such that

$$U_i \cap \Omega = U_i \cap \{(y', y_n) \in \mathbf{R}^{n-1} \times \mathbf{R} : y_n > f(y')\}.$$

Under the uniformly Lipschitz assumption on  $\Omega$ , the extension operator is defined in  $W^{1,p}(\Omega)$ , for all  $p \geq 1$ . We also note that when  $\Omega$  is a bounded domain the above definition reduces to  $\Omega$  being Lipschitz.

In the sequel we set  $a = 1 - 2s$ . Since  $0 < s < 1$  we also have  $-1 < a < 1$ . We first establish the following useful identity:

**Lemma 1.** *Suppose that  $a \in (-1, 1)$  and let  $u \in C_0^\infty(\Omega \times \mathbf{R})$  and  $\phi \in C^2(\Omega \times (0, \infty)) \cap C(\bar{\Omega} \times [0, \infty))$  is such that  $\phi(x, y) > 0$  in  $\Omega \times [0, \infty)$ ,  $\phi(x, y) = 0$  in  $\partial\Omega \times (0, \infty)$ ,*

$$\left| y^a \frac{\phi_y(x, y)}{\phi(x, y)} \right| \leq V(x), \quad y \in (0, 1), \quad x \in \Omega, \quad 0 \leq V(x) \in L^1_{\text{loc}}(\Omega),$$

and for almost everywhere  $x \in \Omega$ , the following limit exists:

$$\lim_{y \rightarrow 0^+} \left( y^a \frac{\phi_y(x, y)}{\phi(x, y)} \right).$$

We also require that the following integrals be finite

$$\int_0^{+\infty} \int_\Omega y^a \frac{|\nabla\phi|^2}{\phi^2} u^2 \, dx \, dy, \quad \int_0^{+\infty} \int_\Omega \frac{|\text{div}(y^a \nabla\phi)|}{\phi} u^2 \, dx \, dy.$$

We then have the identity:

$$\int_0^{+\infty} \int_{\Omega} y^a |\nabla u|^2 \, dx \, dy = \int_0^{+\infty} \int_{\Omega} y^a \left| \nabla u - \frac{\nabla \phi}{\phi} u \right|^2 \, dx \, dy - \int_0^{+\infty} \int_{\Omega} \frac{\operatorname{div}(y^a \nabla \phi)}{\phi} u^2 \, dx \, dy - \int_{\Omega} \lim_{y \rightarrow 0^+} \left( y^a \frac{\phi_y}{\phi} \right) u^2(x, 0) \, dx. \quad (2.1)$$

**Proof.** Expanding the square and integrating by parts, we compute for  $\varepsilon > 0$ ,

$$\begin{aligned} & \int_{\varepsilon}^{\infty} \int_{\Omega} y^a \left| \nabla u - \frac{\nabla \phi}{\phi} u \right|^2 \, dx \, dy \\ &= \int_{\varepsilon}^{\infty} \int_{\Omega} y^a \left( |\nabla u|^2 + \frac{|\nabla \phi|^2}{\phi^2} u^2 - \frac{\nabla \phi}{\phi} \nabla u^2 \right) \, dx \, dy \\ &= \int_{\varepsilon}^{\infty} \int_{\Omega} y^a |\nabla u|^2 \, dx \, dy + \int_{\varepsilon}^{\infty} \int_{\Omega} \frac{\operatorname{div}(y^a \nabla \phi)}{\phi} u^2 \, dx \, dy \\ & \quad + \int_{\Omega} \varepsilon^a \frac{\phi_y(x, \varepsilon)}{\phi(x, \varepsilon)} u^2(x, \varepsilon) \, dx. \end{aligned}$$

We then pass to limit  $\varepsilon \rightarrow 0$  and the result follows easily.  $\square$

We will use Lemma 1 with the following choice:  $\phi(x, y) = d^{-\frac{a}{2}}(x)A(\frac{y}{d(x)})$  for  $y > 0, x \in \Omega$ . The function  $A$  solves the following boundary value problem

$$(t^3 + t)A'' + (a + t^2(2 + a))A' + \frac{(2 + a)a}{4}tA = 0, \quad t > 0, \quad (2.2)$$

with

$$A(0) = 1, \quad \lim_{t \rightarrow +\infty} A(t) = 0. \quad (2.3)$$

Equation (2.2) can also be written in divergence form as

$$(t^a(1 + t^2)A')' + \frac{(2 + a)a}{4}t^a A = 0. \quad (2.4)$$

From now on we will use the following notation:

$$f \sim g, \quad \text{in } U,$$

whenever there exist positive constants  $c_1, c_2$ , such that

$$c_1 g \leq f \leq c_2 g, \quad \text{in } U.$$

We then have the following

**Proposition 7.** *Suppose that  $a \in (-1, 1)$ . The boundary value problem (2.2), (2.3) has a positive decreasing solution  $A$  with the following properties:*

(i) *There exists a positive constant  $\bar{d}_s$  such that*

$$\lim_{t \rightarrow 0^+} t^a A'(t) = -\bar{d}_s,$$

with

$$\bar{d}_s = \frac{(1-a)\Gamma\left(\frac{1+a}{2}\right)\Gamma^2\left(\frac{4-a}{4}\right)}{\Gamma^2\left(\frac{2+a}{4}\right)\Gamma\left(\frac{3-a}{2}\right)} = \frac{2s\Gamma(1-s)\Gamma^2\left(\frac{3+2s}{4}\right)}{\Gamma^2\left(\frac{3-2s}{4}\right)\Gamma(1+s)}.$$

(ii) *For all  $t > 0$ ,*

$$\begin{aligned} A(t) &\sim (1+t^2)^{-\frac{2+a}{4}}, \\ A'(t) &\sim -t^{-a}(1+t^2)^{-\frac{4-a}{4}}. \end{aligned}$$

Moreover,

$$\lim_{t \rightarrow +\infty} \frac{tA'(t)}{A(t)} = -\frac{2+a}{2}.$$

(iii) *There holds:*

$$\bar{d}_s = \int_0^\infty t^a(1+t^2)(A')^2 dt - \frac{(2+a)a}{4} \int_0^\infty t^a A^2 dt, \quad (2.5)$$

(iv) *In case  $a \in (-1, 0]$ , we have*

$$tA'(t) + \frac{a}{2}A(t) \leq 0.$$

Moreover for  $a \in (-1, 0)$  and all  $t > 0$  we have

$$tA'(t) + \frac{a}{2}A(t) \sim -A(t).$$

**Proof.** We change variables in (2.2) by  $z = -t^2$  and define  $B(z)$  such that  $A(t) = B(-t^2)$ , whence  $A_t = -2tB_z$  and  $A_{tt} = -2B_z + 4t^2B_{zz}$ . It then follows that  $B(z)$  satisfies the Gauss hypergeometric equation

$$z(1-z)B'' + \left(\frac{1+a}{2} - \frac{3+a}{2}z\right)B' - \frac{a(2+a)}{16}B = 0, \quad -\infty < z < 0,$$

whose general solution is given by

$$B(z) = C_1 F_1\left(\frac{a}{4}, \frac{2+a}{4}, \frac{1+a}{2}; z\right) + C_2 z^{\frac{1-a}{2}} F_2\left(\frac{2-a}{4}, \frac{4-a}{4}, \frac{3-a}{2}; z\right);$$

see [1, Section 15.5 as well as 15.1] for the definition and basic properties of the function  $F$ . It follows that

$$\begin{aligned} A(t) &= C_1 F_1\left(\frac{a}{4}, \frac{2+a}{4}, \frac{1+a}{2}; -t^2\right) \\ &\quad + C_2 t^{1-a} e^{\frac{i\pi(1-a)}{2}} F_2\left(\frac{2-a}{4}, \frac{4-a}{4}, \frac{3-a}{2}; -t^2\right). \end{aligned} \quad (2.6)$$

Since  $F(\alpha, \beta, \gamma; 0) = 1$  for any  $\alpha, \beta, \gamma$ , the condition  $A(0) = 1$  implies that  $C_1 = 1$ . We then have

$$\begin{aligned} \bar{d}_s &= - \lim_{t \rightarrow 0^+} t^a A'(t) \\ &= - \lim_{t \rightarrow 0^+} t^a (-2tF_1' + (1-a)C_2 e^{\frac{i\pi(1-a)}{2}} t^{-a} F_2 - 2C_2 t^{2-a} e^{\frac{i\pi(1-a)}{2}} F_2') \\ &= -(1-a)C_2 e^{\frac{i\pi(1-a)}{2}}. \end{aligned} \tag{2.7}$$

In the above calculation we have also used the fact that

$$F'(\alpha, \beta, \gamma; z) = \frac{d}{dz} F(\alpha, \beta, \gamma; z) = \frac{\alpha\beta}{\gamma} F(\alpha + 1, \beta + 1, \gamma + 1; z).$$

We next compute the behavior of  $A$  at infinity. To this end we will use the inversion formula, valid for any  $\alpha, \beta, \gamma$  and  $|\arg(-z)| < \pi$ :

$$\begin{aligned} F(\alpha, \beta, \gamma; z) &= \frac{\Gamma(\gamma)\Gamma(\beta-\alpha)}{\Gamma(\beta)\Gamma(\gamma-\alpha)} (-z)^{-\alpha} F\left(\alpha, 1-\gamma+\alpha, 1-\beta+\alpha; \frac{1}{z}\right) \\ &\quad + \frac{\Gamma(\gamma)\Gamma(\alpha-\beta)}{\Gamma(\alpha)\Gamma(\gamma-\beta)} (-z)^{-\beta} F\left(\beta, 1-\gamma+\beta, 1-\alpha+\beta; \frac{1}{z}\right). \end{aligned}$$

We then calculate

$$\lim_{t \rightarrow +\infty} t^{\frac{a}{2}} A(t) = \frac{\Gamma\left(\frac{1+a}{2}\right)\Gamma\left(\frac{1}{2}\right)}{\Gamma^2\left(\frac{2+a}{4}\right)} + C_2 e^{\frac{i\pi(1-a)}{2}} \frac{\Gamma\left(\frac{3-a}{2}\right)\Gamma\left(\frac{1}{2}\right)}{\Gamma^2\left(\frac{4-a}{4}\right)}.$$

To make this limit equal to zero, we choose

$$C_2 = -e^{-\frac{i\pi(1-a)}{2}} \frac{\Gamma\left(\frac{1+a}{2}\right)\Gamma^2\left(\frac{4-a}{4}\right)}{\Gamma^2\left(\frac{2+a}{4}\right)\Gamma\left(\frac{3-a}{2}\right)}.$$

Combining this with (2.7) we conclude

$$\bar{d}_s = \frac{(1-a)\Gamma\left(\frac{1+a}{2}\right)\Gamma^2\left(\frac{4-a}{4}\right)}{\Gamma^2\left(\frac{2+a}{4}\right)\Gamma\left(\frac{3-a}{2}\right)} = \frac{2s\Gamma(1-s)\Gamma^2\left(\frac{3+2s}{4}\right)}{\Gamma^2\left(\frac{3-2s}{4}\right)\Gamma(1+s)}. \tag{2.8}$$

At this point, both constants  $C_1, C_2$ , in (2.6) have been identified. After some lengthy but straightforward calculations, we find that as  $t \rightarrow +\infty$ ,

$$A(t) \sim t^{-\frac{2+a}{2}}, \quad A'(t) \sim t^{-\frac{4+a}{2}}. \tag{2.9}$$

In addition, we get

$$\lim_{t \rightarrow +\infty} \frac{tA'(t)}{A(t)} = -\frac{2+a}{2}.$$

Using (2.4) and the above asymptotics, we easily conclude that the solution  $A$  is energetic, that is,

$$\int_0^\infty t^a (1+t^2)(A')^2 dt + \int_0^\infty t^a A^2 dt < \infty.$$

Multiplying (2.4) by  $A$  and integrating by parts in  $(0, \infty)$  we arrive at (2.5)

To prove the positivity and monotonicity of  $A$ , we next change variables by:

$$B(s) = (1 + t^2)^{\frac{a}{4}} A(t), \quad s = 1/t.$$

It follows that  $B$  satisfies the equation

$$(1 + s^2)^2 B'' + (2 - a)s(1 + s^2)B' - \frac{a^2}{4}B = 0, \quad s \in (0, +\infty),$$

with  $B(0) = 0$  and  $B(+\infty) = 1$ . A standard maximum principle argument shows that  $B$  is positive. Consequently,  $A$  is positive and the monotonicity of  $A$  follows easily.

The positivity and monotonicity of  $A$  in connection with the asymptotics of  $A$  easily yield part (ii) of the proposition.

Part (iv) follows easily from the monotonicity of  $A$  and part (ii).  $\square$

Using the asymptotics of  $A(t)$ , from the previous Proposition we easily obtain the following uniform asymptotics for  $\phi$

**Lemma 2.** *Suppose  $a \in (-1, 1)$  and let  $\phi$  be given by*

$$\phi(x, y) = d^{-\frac{a}{2}}(x)A\left(\frac{y}{d(x)}\right), \quad y > 0, \quad x \in \Omega \subset \mathbf{R}^n,$$

where  $A$  solves (2.2), (2.3).

(i) Then

$$\phi(x, y) \sim \frac{d}{(d^2 + y^2)^{\frac{2+a}{4}}}, \quad y > 0, \quad x \in \Omega.$$

Concerning the gradient of  $\phi$ , for  $a \in (-1, 0]$ , we have

$$|\nabla_{(x,y)}\phi(x, y)| \sim \frac{1}{(d^2 + y^2)^{\frac{2+a}{4}}}, \quad y > 0, \quad x \in \Omega,$$

whereas for  $a \in (0, 1)$ ,

$$|\nabla_{(x,y)}\phi(x, y)| \sim \frac{y^{-a}}{(d^2 + y^2)^{\frac{2-a}{4}}}, \quad y > 0, \quad x \in \Omega.$$

(ii) If  $\Omega$  satisfies  $-\Delta d(x) \geq 0$  for  $x \in \Omega$ , then for  $a \in (-1, 0)$

$$-\operatorname{div}(y^a \nabla \phi)\phi \sim \frac{y^a}{(d^2 + y^2)^{\frac{2+a}{2}}}(-d \Delta d), \quad y > 0, \quad x \in \Omega,$$

whereas for  $a = 0$ ,

$$-\operatorname{div}(\nabla \phi)\phi \sim \frac{y}{(d^2 + y^2)^{\frac{3}{2}}}(-d \Delta d), \quad y > 0, \quad x \in \Omega.$$

We are now ready to give the proof of Theorem 1.

**Proof of Theorem 1 parts (i) and (ii).** We assume that  $s \in [\frac{1}{2}, 1)$  or, equivalently,  $a \in (-1, 0]$ . We will use Lemma 1 with the test function  $\phi$  given by

$$\phi(x, y) = d^{-\frac{a}{2}}(x)A\left(\frac{y}{d(x)}\right), \quad y > 0, \quad x \in \Omega \subset \mathbf{R}^n,$$

where  $A$  solves (2.2), (2.3). Using Proposition 7 and Lemma 2 we see that all hypotheses of Lemma 1 are satisfied. In particular, for  $t = \frac{y}{d}$  we compute, for  $x \in \Omega$ ,

$$\begin{aligned} -\lim_{y \rightarrow 0^+} \left( y^a \frac{\phi_y}{\phi} \right) &= -\lim_{y \rightarrow 0^+} \left( t^a \frac{A'(t)}{d^{1-a}A(t)} \right) = \frac{1}{d^{1-a}(x)} \lim_{t \rightarrow 0^+} \left( -\frac{t^a A'(t)}{A(t)} \right) \\ &= \frac{\bar{d}_s}{d^{1-a}(x)}. \end{aligned} \tag{2.10}$$

We also have

$$\begin{aligned} -\operatorname{div}(y^a \nabla \phi) &= -y^{a-1} d^{-1-\frac{a}{2}} \left[ (t^3 + t)A'' + (a + t^2(2+a))A' + \frac{(2+a)a}{4}tA \right] \\ &\quad - y^{a-1} d^{-1-\frac{a}{2}} \left[ (-d\Delta d) \left( t^2 A' + \frac{at}{2} A \right) \right] \\ &= -y^{a-1} d^{-1-\frac{a}{2}} \left[ (-d\Delta d) \left( t^2 A' + \frac{at}{2} A \right) \right], \end{aligned}$$

therefore,

$$-\operatorname{div}(y^a \nabla \phi) \geq 0, \quad x \in \Omega, \quad y > 0.$$

From Lemma 1 we get

$$\begin{aligned} \int_0^{+\infty} \int_{\Omega} y^a |\nabla u|^2 \, dx \, dy &\geq \bar{d}_s \int_{\Omega} \frac{u^2(x, 0)}{d^{1-a}(x)} \, dx + \int_0^{+\infty} \int_{\Omega} y^a \left| \nabla u - \frac{\nabla \phi}{\phi} u \right|^2 \, dx \, dy \\ &\quad - \int_0^{+\infty} \int_{\Omega} \frac{\operatorname{div}(y^a \nabla \phi)}{\phi} u^2 \, dx \, dy, \end{aligned} \tag{2.11}$$

from which the trace Hardy inequality follows directly. This relation will be used later on, in Sections 5 and 6, to obtain the Sobolev term as well.

We continue with the proof of the optimality of the Hardy constant  $\bar{d}_s$ . Let

$$Q[u] := \frac{\int_0^{+\infty} \int_{\Omega} y^a |\nabla u|^2 \, dx \, dy}{\int_{\Omega} \frac{u^2(x, 0)}{d^{1-a}(x)} \, dx} =: \frac{N[u]}{D[u]}. \tag{2.12}$$

We have that  $Q[u] \geq \bar{d}_s$ . Here we will show that there exists a sequence of functions  $u_\varepsilon$  such that  $\lim_{\varepsilon \rightarrow 0} Q[u_\varepsilon] = \bar{d}_s$ , and therefore  $\bar{d}_s$  is the best constant.

We first assume for simplicity that the boundary of  $\Omega$  is flat in a neighborhood  $V$  of a point  $x_0 \in \partial\Omega$ . The neighborhood of the point  $x_0$  is assumed to contain a ball centered at  $x_0$  with radius, say,  $3\delta$ . Locally around  $x_0$  the boundary is given by  $x_n = 0$ , whereas the interior of  $\Omega$  corresponds to  $x_n > 0$ . We also write  $x = (x', x_n)$ . Clearly, for  $x \in \Omega \cap V$  we have that  $d(x) = x_n$ .



We next define two suitable cutoff functions. Let  $\psi(x') \in C_0^\infty(B_\delta)$ , where  $B_\delta \subset \partial\Omega \subset \mathbf{R}^{n-1}$  is the ball centered at  $x_0$  with radius  $\delta$ . Also, the nonnegative function  $h(x_n) \in C^\infty(\mathbf{R}^+)$  is such that  $h(x_n) = 0$  for  $x_n \geq 2\delta$  and  $h(x_n) = 1$  for  $0 \leq x_n \leq \delta$ . We will use the following test function:

$$u_\varepsilon(x', x_n, y) = \begin{cases} h(x_n)\psi(x')x_n^{-\frac{a}{2}}A\left(\frac{y}{x_n}\right), & y \geq \varepsilon \\ h(x_n)\psi(x')x_n^{-\frac{a}{2}}A\left(\frac{\varepsilon}{x_n}\right), & 0 \leq y < \varepsilon. \end{cases} \quad (2.13)$$

We have that

$$Q[u_\varepsilon] = \frac{\int_0^{+\infty} dy \int_0^{2\delta} dx_n \int_{B_\delta} dx' y^a |\nabla u_\varepsilon|^2}{\int_0^{2\delta} dx_n \int_{B_\delta} dx' \frac{u_\varepsilon^2}{x_n^{1-a}}} = \frac{N[u_\varepsilon]}{D[u_\varepsilon]}. \quad (2.14)$$

Concerning the denominator, we compute

$$\begin{aligned} D[u_\varepsilon] &= \int_{B_\delta} \psi^2(x') dx' \int_0^\delta x_n^{-1} A^2\left(\frac{\varepsilon}{x_n}\right) dx_n + O_\varepsilon(1) \\ &= \int_{B_\delta} \psi^2(x') dx' \int_{\varepsilon/\delta}^{+\infty} \frac{A^2(t)}{t} dt + O_\varepsilon(1). \end{aligned} \quad (2.15)$$

We next calculate the numerator. At first, we break  $N$  into two pieces:

$$N[u_\varepsilon] = \int_0^\varepsilon dy + \int_\varepsilon^{+\infty} dy =: N_1[u_\varepsilon] + N_2[u_\varepsilon].$$

Using the specific form of  $u_\varepsilon$  and elementary estimates, we calculate:

$$\begin{aligned} N_2[u_\varepsilon] &= \int_{B_\delta} \psi^2(x') dx' \int_\varepsilon^{+\infty} dy \int_0^\delta dx_n \frac{y^a}{x_n^{a+2}} \\ &\quad \times \left[ \left( -\frac{a}{2} A\left(\frac{y}{x_n}\right) - \frac{y}{x_n} A'\left(\frac{y}{x_n}\right) \right)^2 + A'^2\left(\frac{y}{x_n}\right) \right] \\ &\quad + \int_{B_\delta} |\nabla \psi(x')|^2 dx' \int_\varepsilon^{+\infty} dy \int_0^\delta dx_n y^a x_n^{-a} A^2\left(\frac{y}{x_n}\right) + O_\varepsilon(1) \\ &=: N_{21}[u_\varepsilon] + N_{22}[u_\varepsilon] + O_\varepsilon(1). \end{aligned}$$

We note that as  $\varepsilon \rightarrow 0$ ,

$$\begin{aligned} N_{22}[u_\varepsilon] &= \int_{B_\delta} |\nabla \psi(x')|^2 dx' \int_0^\delta x_n \int_{\varepsilon/x_n}^{+\infty} t^a A^2(t) dt dx_n \\ &= O_\varepsilon(1). \end{aligned}$$

Concerning  $N_{21}[u_\varepsilon]$ , changing variables by  $t = \frac{y}{x_n}$  we write:

$$\begin{aligned} N_{21}[u_\varepsilon] &= \int_{B_\delta} \psi^2(x') dx' \int_\varepsilon^{+\infty} \frac{dy}{y} \int_{y/\delta}^{+\infty} t^a \left[ A'^2(t) + \left( \frac{a}{2} A(t) + t A'(t) \right)^2 \right] dt \\ &= \int_{B_\delta} \psi^2(x') dx' \int_\varepsilon^{+\infty} \frac{dy}{y} \int_{y/\delta}^{+\infty} t^a \left[ (1+t^2)A'^2 + atAA' + \frac{a^2}{4}A^2 \right] dt. \end{aligned}$$

Integrating by parts the term containing the factors  $AA'$ , then using the equation satisfied by  $A$  (see (2.4)), we get

$$\begin{aligned} & \int_{y/\delta}^{+\infty} \left[ t^a(1+t^2)A'^2 + at^{1+a}AA' + \frac{a^2}{4}t^aA^2 \right] dt \\ &= \int_{y/\delta}^{+\infty} \left[ t^a(1+t^2)A'^2 - \frac{a(2+a)}{4}t^aA^2 \right] dt + \frac{1}{2}at^{1+a}A^2(t) \Big|_{t=y/\delta} \\ &= -t^a(1+t^2)A(t)A'(t) \Big|_{t=y/\delta} + \frac{1}{2}at^{1+a}A^2(t) \Big|_{t=y/\delta}, \end{aligned}$$

whence,

$$N_{21}[u_\varepsilon] = - \int_{B_\delta} \psi^2(x') dx' \int_{\varepsilon/\delta}^{+\infty} \frac{1}{t} t^a(1+t^2)A(t)A'(t) dt + O_\varepsilon(1).$$

It is not difficult to show that  $N_1[u_\varepsilon] = O_\varepsilon(1)$ , and therefore  $N[u_\varepsilon] = N_{21}[u_\varepsilon] + O_\varepsilon(1)$ . Using (2.15), in addition, we can form the quotient

$$\begin{aligned} \lim_{\varepsilon \rightarrow 0} Q[u_\varepsilon] &= \lim_{\varepsilon \rightarrow 0} \frac{- \int_{B_\delta} \psi^2(x') dx' \int_{\varepsilon/\delta}^{+\infty} \frac{1}{t} t^a(1+t^2)A(t)A'(t) dt + O_\varepsilon(1)}{\int_{B_\delta} \psi^2(x') dx' \int_{\varepsilon/\delta}^{+\infty} \frac{A^2(t)}{t} dt + O_\varepsilon(1)} \\ &= \lim_{\varepsilon \rightarrow 0} \frac{- \int_{\varepsilon/\delta}^{+\infty} \frac{1}{t} t^a(1+t^2)A(t)A'(t) dt}{\int_{\varepsilon/\delta}^{+\infty} \frac{A^2(t)}{t} dt} \\ &= - \lim_{\sigma \rightarrow 0} \frac{\sigma^a(1+\sigma^2)A'(\sigma)}{A(\sigma)} \\ &= \bar{d}_s, \end{aligned} \tag{2.16}$$

where we used L'Hopital's rule and then part (i) of Proposition 7.

Let us now consider the general case. We assume that  $\partial\Omega$  is  $C^1$  in a neighborhood of a point  $\bar{x}_0$ , which we take to be the origin  $0 \in \partial\Omega$ . Thus, locally,  $\partial\Omega$  is the graph of a function  $\bar{x}_n = \gamma(\bar{x}')$ , with  $\gamma(0) = 0$  and  $\nabla\gamma(0) = 0$ . We also assume that the interior of  $\Omega$  corresponds to  $\bar{x}_n > \gamma(\bar{x}')$ . Then the following change of coordinates straightens the boundary in a neighborhood of the origin:  $x_i = \bar{x}_i, i = 1, 2, \dots, n-1$ , and  $x_n = \bar{x}_n - \gamma(\bar{x}')$ ; see for example [22, Appendix C]. We assume that inside the ball  $B(0, 3\delta)$  (in the  $x$ -space) the image of  $\partial\Omega$  is flat. We then consider the test function  $v_\varepsilon(\bar{x}, y) = u_\varepsilon(x, y)$ . Clearly  $v_\varepsilon(\bar{x}, y)$  is zero away from a neighborhood of the origin, say  $U$ , and elementary calculations show that

$$\nabla_{\bar{x}} v_\varepsilon = \nabla_x u_\varepsilon - u_{\varepsilon, x_n} \nabla_{\bar{x}} \gamma(\bar{x}'),$$

whence,

$$|\nabla_{\bar{x}} v_\varepsilon - \nabla_x u_\varepsilon| \leq |\nabla_{\bar{x}} \gamma(\bar{x}')| |\nabla_x u_\varepsilon| = o_\delta(1) |\nabla_x u_\varepsilon|.$$

It then follows that

$$|\nabla_{\bar{x}} v_\varepsilon| = |\nabla_x u_\varepsilon| (1 + o_\delta(1)).$$

On the other hand, for  $\bar{x} \in U$  and  $d(\bar{x}) = \text{dist}(\bar{x}, \partial\Omega)$ , we have that

$$d(\bar{x}) = (\bar{x}_n - \gamma(\bar{x}'))(1 + |\nabla_{\bar{x}}\gamma(\bar{x}')|^2)^{1/2} = x_n(1 + o_\delta(1)).$$

We finally note that the Jacobian of the above transformation is one, and therefore  $dx = d\bar{x}$ . We then compute

$$Q[v_\varepsilon(\bar{x}, y)] = Q[u_\varepsilon(x, y)](1 + o_\delta(1)),$$

where  $Q[u_\varepsilon(x, y)]$  is given in (2.14). Since  $\delta$  can be taken as small as we like, the result follows easily, using the calculations from the flat case.  $\square$

### 3. The Trace Hardy Inequality II

In this section we will prove the trace Hardy inequality contained in Theorem 4. We first establish the analogue of Lemma 1:

**Lemma 3.** *Suppose that  $a \in (-1, 1)$  and let  $u \in C_0^\infty(\mathbf{R}^n \times \mathbf{R})$  such that  $u(\cdot, 0) \in C_0^\infty(\Omega)$ . Let  $\phi \in C^2(\mathbf{R}^n \times (0, \infty)) \cap C(\mathbf{R}^n \times [0, \infty))$  such that  $\phi(x, y) > 0$  in  $\mathbf{R}^n \times [0, \infty)$ ,  $\phi(x, 0) = 0$  in  $x \in \mathcal{C}\Omega$ ,*

$$|y^a \frac{\phi_y(x, y)}{\phi(x, y)}| \leq V(x), \quad y \in (0, 1), \quad x \in \mathbf{R}^n, \quad 0 \leq V(x) \in L_{\text{loc}}^1(\mathbf{R}^n).$$

Moreover for almost everywhere  $x \in \Omega$ , the following limit exists:

$$\lim_{y \rightarrow 0^+} \left( y^a \frac{\phi_y(x, y)}{\phi(x, y)} \right).$$

We also require that the following integrals be finite

$$\int_0^{+\infty} \int_{\mathbf{R}^n} y^a \frac{|\nabla\phi|^2}{\phi^2} u^2 \, dx \, dy, \quad \int_0^{+\infty} \int_{\mathbf{R}^n} \frac{|\text{div}(y^a \nabla\phi)|}{\phi} u^2 \, dx \, dy.$$

We then have the identity

$$\begin{aligned} \int_0^{+\infty} \int_{\mathbf{R}^n} y^a |\nabla u|^2 \, dx \, dy &= \int_0^{+\infty} \int_{\mathbf{R}^n} y^a \left| \nabla u - \frac{\nabla\phi}{\phi} u \right|^2 \, dx \, dy \\ &- \int_0^{+\infty} \int_{\mathbf{R}^n} \frac{\text{div}(y^a \nabla\phi)}{\phi} u^2 \, dx \, dy - \int_{\Omega} \lim_{y \rightarrow 0^+} \left( y^a \frac{\phi_y}{\phi} \right) u^2(x, 0) \, dx. \end{aligned} \quad (3.1)$$

The proof of this lemma is quite similar to the proof of Lemma 1, so we omit it.

This time we will choose the test function to be of the form

$$\phi(x, y) = \begin{cases} (y^2 + d^2)^{-\frac{a}{4}} B\left(\frac{d}{y}\right), & x \in \Omega, \quad y > 0 \\ (y^2 + d^2)^{-\frac{a}{4}} B\left(-\frac{d}{y}\right), & x \in \mathcal{C}\Omega, \quad y > 0, \end{cases} \quad (3.2)$$

where function  $B$  is the solution of the boundary value problem

$$(1 + t^2)^2 B'' + (2 - a)t(1 + t^2)B' - \frac{a^2}{4}B = 0, \quad t \in (-\infty, +\infty), \quad (3.3)$$

complemented with the conditions

$$B(-\infty) = 0, \quad B(+\infty) = 1. \quad (3.4)$$

We note that this can be written in divergence form as

$$((1 + t^2)^{1-\frac{a}{2}} B'(t))' - \frac{a^2}{4}(1 + t^2)^{-1-\frac{a}{2}} B(t) = 0, \quad t \in \mathbf{R}. \quad (3.5)$$

We next collect some properties of  $B$  that will be used later on.

**Proposition 8.** *Suppose that  $a \in (-1, 1)$ . The boundary value problem (3.3), (3.4) has a positive increasing solution  $B$  with the following properties:*

(i) *There exists a positive constant  $\bar{k}_s$  such that*

$$\lim_{t \rightarrow +\infty} (1 + t^2)^{\frac{2-a}{2}} B'(t) =: \bar{k}_s, \quad (3.6)$$

where

$$\bar{k}_s = \frac{2^a \Gamma^2\left(\frac{2-a}{2}\right) \Gamma\left(\frac{1+a}{2}\right)}{\pi \Gamma\left(\frac{1-a}{2}\right)} = \frac{2^{1-2s} \Gamma^2\left(s + \frac{1}{2}\right) \Gamma(1-s)}{\pi \Gamma(s)}.$$

(ii) *We have*

$$\begin{aligned} B(t) &\sim 1, \quad t > 0 \\ B(t) &\sim (1 + t^2)^{-\frac{1-a}{2}} \quad t < 0, \\ B'(t) &\sim (1 + t^2)^{-\frac{2-a}{2}} \quad t \in \mathbf{R}. \end{aligned}$$

(iii) *There holds:*

$$\bar{k}_s = \int_{-\infty}^{+\infty} \left[ (1 + t^2)^{1-\frac{a}{2}} B'^2(t) + \frac{a^2}{4}(1 + t^2)^{-1-\frac{a}{2}} B^2(t) \right] dt.$$

(iv) *In the case where  $a \in (-1, 0]$ , we have*

$$(1 + t^2)B'(t) - \frac{a}{2}tB(t) > 0, \quad t \in \mathbf{R}.$$

Moreover, for  $a \in (-1, 0)$

$$(1 + t^2)B'(t) - \frac{a}{2}tB(t) \sim (1 + t^2)^{\frac{1}{2}}, \quad t > 0.$$

**Proof.** When  $a = 0$  the ODE can be easily solved by a straightforward integration. For the general case, we first change variables by  $B(t) = (1 + t^2)^{\frac{a}{4}} f(t)$  to obtain

$$(1 + t^2)f'' + 2tf' + \frac{a(2 - a)}{4}f = 0.$$

We next change variables by  $g(z) = f(t)$ ,  $z = it$ , so that  $g$  satisfies the equation

$$(1 - z^2)g'' - 2zg' + \nu(\nu + 1)g = 0, \quad \nu = -\frac{a}{2}. \tag{3.7}$$

The solution of this is given in [1, Section 8.1]:

$$g(z) = \begin{cases} C_1^+ P_\nu(z) + C_2^+ Q_\nu(z), & \text{Im } z > 0, \\ C_1^- P_\nu(z) + C_2^- Q_\nu(z), & \text{Im } z < 0. \end{cases} \tag{3.8}$$

We also have that

$$B(t) = (1 + t^2)^{-\frac{\nu}{2}}g(it).$$

The conditions at infinity then become

$$\lim_{t \rightarrow +\infty} t^{-\nu}g(it) = 1, \quad \lim_{t \rightarrow -\infty} (-t)^{-\nu}g(it) = 0. \tag{3.9}$$

To find the constants in (3.8) we will satisfy the conditions at infinity (3.9) and we will match both  $g$  and  $g'$  at  $z = 0$ . That is, we will ask

$$g(+i0) = g(-i0), \quad g'(+i0) = g'(-i0). \tag{3.10}$$

We recall from [1, Section 8.1] that for  $|z| > 1$ :

$$\begin{aligned} P_\nu(z) &= \Delta_1 z^{-\nu-1} F\left(\frac{\nu+1}{2}, \frac{\nu+2}{2}, \frac{2\nu+3}{2}; \frac{1}{z^2}\right) \\ &\quad + \Delta_2 z^\nu F\left(\frac{-\nu}{2}, \frac{1-\nu}{2}, \frac{1-2\nu}{2}; \frac{1}{z^2}\right), \\ Q_\nu(z) &= E_1 z^{-\nu-1} F\left(\frac{\nu+2}{2}, \frac{\nu+1}{2}, \frac{2\nu+3}{2}; \frac{1}{z^2}\right), \end{aligned}$$

where,

$$\begin{aligned} \Delta_1 &= \frac{2^{-\nu-1}\pi^{-\frac{1}{2}}\Gamma(-\nu - \frac{1}{2})}{\Gamma(-\nu)}, & \Delta_2 &= \frac{2^\nu\pi^{-\frac{1}{2}}\Gamma(\nu + \frac{1}{2})}{\Gamma(1 + \nu)}, \\ E_1 &= \frac{2^{-\nu-1}\pi^{\frac{1}{2}}\Gamma(1 + \nu)}{\Gamma(\frac{3}{2} + \nu)}. \end{aligned}$$

From the asymptotics when  $t \rightarrow \pm\infty$ , we easily conclude that

$$C_1^+ = \frac{i^{-\nu}}{\Delta_2}, \quad C_1^- = 0. \tag{3.11}$$

We next see what happens near zero. For  $|z| < 1$  we have that

$$P_\nu(z) = B_1 F\left(-\frac{\nu}{2}, \frac{\nu+1}{2}, \frac{1}{2}; z^2\right) + B_2 z F\left(\frac{1-\nu}{2}, \frac{2+\nu}{2}, \frac{3}{2}; z^2\right),$$

$$Q_\nu^\pm(z) = \Gamma_1 e^{\pm \frac{i\pi}{2}(-\nu-1)} F\left(-\frac{\nu}{2}, \frac{\nu+1}{2}, \frac{1}{2}; z^2\right) + \Gamma_2 e^{\pm \frac{i\pi}{2}(-\nu)} z F\left(\frac{1-\nu}{2}, \frac{\nu+2}{2}, \frac{3}{2}; z^2\right),$$

where the plus sign corresponds to  $\text{Im } z > 0$  and the minus sign to  $\text{Im } z < 0$ . The values of the constants are given by:

$$B_1 = \frac{\pi^{\frac{1}{2}}}{\Gamma\left(\frac{1-\nu}{2}\right)\Gamma\left(\frac{2+\nu}{2}\right)}, \quad B_2 = \frac{-2\pi^{\frac{1}{2}}}{\Gamma\left(\frac{1+\nu}{2}\right)\Gamma\left(\frac{-\nu}{2}\right)},$$

$$\Gamma_1 = \frac{\pi^{\frac{1}{2}}\Gamma\left(\frac{1+\nu}{2}\right)}{2\Gamma\left(1+\frac{\nu}{2}\right)}, \quad \Gamma_2 = \frac{\pi^{\frac{1}{2}}\Gamma\left(1+\frac{\nu}{2}\right)}{\Gamma\left(\frac{1+\nu}{2}\right)}.$$

An easy calculation shows that the matching condition (3.10) yields

$$C_2^- \Gamma_1 e^{\frac{i\pi}{2}(\nu+1)} = C_1^+ B_1 + C_2^+ \Gamma_1 e^{\frac{i\pi}{2}(-\nu-1)},$$

$$C_2^- \Gamma_2 e^{\frac{i\pi}{2}\nu} = C_1^+ B_2 + C_2^+ \Gamma_2 e^{\frac{i\pi}{2}(-\nu)},$$

from which it follows that

$$C_2^+ = -\frac{C_1^+}{2} e^{\frac{i\pi}{2}\nu} \left[ \frac{B_2}{\Gamma_2} + i \frac{B_1}{\Gamma_1} \right]$$

$$C_2^- = \frac{C_1^+}{2} e^{-\frac{i\pi}{2}\nu} \left[ \frac{B_2}{\Gamma_2} - i \frac{B_1}{\Gamma_1} \right]. \tag{3.12}$$

Thus all constants in (3.8) have been computed (see (3.11) and (3.12)), and therefore  $g(z)$  is now completely known.

The asymptotics of  $g$  for  $|z| \rightarrow +\infty$  are

$$g(z) = C_1^\pm \Delta_2 z^\nu + (C_1^\pm \Delta_1 + C_2^\pm E_1) z^{-\nu-1} + o(|z|^{-\nu-1}),$$

$$g'(z) = C_1^\pm \Delta_2 \nu z^{\nu-1} - (\nu+1)[C_1^\pm \Delta_1 + C_2^\pm E_1] z^{-\nu-2} + O(|z|^{\nu-3}),$$

where the plus sign corresponds to  $\text{Im } z > 0$  and the minus sign to  $\text{Im } z < 0$ . We have that  $B(t) = (1+t^2)^{-\frac{\nu}{2}} g(it)$ , whence we get

$$B(t) = i^{1+\nu} C_2^- E_1 (-t)^{-2\nu-1} + o((-t)^{-2\nu-1}), \quad t \rightarrow -\infty.$$

Concerning the derivative, we have for  $z = it$

$$B'(t) = -\nu t (t^2 + 1)^{-\frac{\nu}{2}-1} g(z) + i (1+t^2)^{-\frac{\nu}{2}} g'(z).$$

Whence,

$$B'(t) = (2\nu+1) i^{1-\nu} (C_1^+ \Delta_1 + C_2^+ E_1) t^{-2\nu-2} + o(t^{-2\nu-2}), \quad t \rightarrow +\infty,$$

$$B'(t) = (2\nu+1) i^{1+\nu} C_2^- E_1 (-t)^{-2\nu-2} + o((-t)^{-2\nu-2}), \quad t \rightarrow -\infty.$$

This completes the proof of part (ii) of the proposition.

We next give the proof of part (i). From (3.6) and the asymptotics of  $B(t)$  for  $t \rightarrow +\infty$ , we compute

$$\bar{k}_s = \frac{(2\nu + 1)}{2} i^{1-2\nu} \frac{E_1}{\Delta_2} \left( 2 \frac{\Delta_1}{E_1} - i^\nu \frac{B_2}{\Gamma_2} - i^{\nu+1} \frac{B_1}{\Gamma_1} \right). \quad (3.13)$$

Using the explicit values of the constants we calculate:

$$\begin{aligned} \frac{E_1}{\Delta_2} &= \frac{2^{-2\nu-1} \pi \Gamma^2(1 + \nu)}{\Gamma\left(\frac{1}{2} + \nu\right) \Gamma\left(\frac{3}{2} + \nu\right)}, & \frac{\Delta_1}{E_1} &= \frac{\sin(\pi\nu)}{\pi \cos(\pi\nu)}, \\ \frac{B_2}{\Gamma_2} &= \frac{2 \sin\left(\frac{\pi\nu}{2}\right)}{\pi}, & \frac{B_1}{\Gamma_1} &= \frac{2 \cos\left(\frac{\pi\nu}{2}\right)}{\pi}. \end{aligned}$$

Plugging these into (3.13) we conclude that [recall that  $\nu = -a/2 = s - 1/2$ ]

$$\begin{aligned} \bar{k}_s &= \frac{2^{-2\nu}}{\pi} \frac{\Gamma^2(1 + \nu) \Gamma\left(\frac{1}{2} - \nu\right)}{\Gamma\left(\frac{1}{2} + \nu\right)} = \frac{2^a \Gamma^2\left(\frac{2-a}{2}\right) \Gamma\left(\frac{1+a}{2}\right)}{\pi \Gamma\left(\frac{1-a}{2}\right)} \\ &= \frac{2^{1-2s}}{\pi} \frac{\Gamma^2\left(s + \frac{1}{2}\right) \Gamma(1 - s)}{\Gamma(s)}. \end{aligned} \quad (3.14)$$

To prove part (iii) we use part (i) and we integrate the ODE (3.5).

By standard maximum principle arguments, the solution  $B(t)$  of (3.3) subject to (3.4) is positive and increasing. To prove part (iv) assuming that  $a \in (-1, 0)$ , we set  $f(t) = (1 + t^2)^{-\frac{a}{4}} B(t)$  so that

$$(1 + t^2) f'' + 2t f' + \frac{a(2 - a)}{4} f = 0,$$

and a similar maximum principle argument shows that  $f(t)$  is also increasing. Since

$$f'(t) = (1 + t^2)^{-\frac{a}{4}-1} \left[ (1 + t^2) B' - \frac{a}{2} t B \right],$$

we conclude that

$$(1 + t^2) B' - \frac{a}{2} t B > 0, \quad t \in \mathbf{R}, \quad a \leq 0.$$

Using the asymptotics of  $B, B'$  from part (ii) we conclude the proof of part (iv). □

Using the asymptotics of  $B(t)$  from the previous proposition, we easily obtain the following uniform asymptotics for  $\phi$ .

**Lemma 4.** *Suppose  $a \in (-1, 1)$  and let  $\phi$  be given by*

$$\phi(x, y) = \begin{cases} (y^2 + d^2)^{-\frac{a}{4}} B\left(\frac{d}{y}\right), & x \in \Omega, \quad y > 0 \\ (y^2 + d^2)^{-\frac{a}{4}} B\left(-\frac{d}{y}\right), & x \in \mathcal{C}\Omega, \quad y > 0, \end{cases}$$

where  $B$  solves (3.3), (3.4).

(i) Then

$$\phi(x, y) \sim \begin{cases} (y^2 + d^2)^{-\frac{a}{4}}, & x \in \Omega, y > 0 \\ y^{1-a}(y^2 + d^2)^{\frac{a-2}{4}}, & x \in \mathcal{C}\Omega, y > 0. \end{cases}$$

Concerning the gradient of  $\phi$ , for  $a \in (-1, 0]$ , we have

$$|\nabla\phi(x, y)| \sim \begin{cases} (y^2 + d^2)^{-\frac{a+2}{4}}, & x \in \Omega, y > 0 \\ y^{-a}(y^2 + d^2)^{\frac{a-2}{4}}, & x \in \mathcal{C}\Omega, y > 0, \end{cases}$$

whereas for  $a \in (0, 1)$

$$|\nabla\phi(x, y)| \sim y^{-a}(y^2 + d^2)^{\frac{a-2}{4}}, \quad x \in \mathbf{R}^n, y > 0.$$

(ii) If  $\Omega$  satisfies  $-\Delta d(x) \geq 0$  for  $x \in \Omega$ , then for  $a \in (-1, 0)$

$$-\operatorname{div}(y^a \nabla\phi)\phi \sim \frac{y^a}{d(d^2 + y^2)^{\frac{1+a}{2}}}(-d\Delta d), \quad y > 0, x \in \Omega,$$

whereas for  $a = 0$ ,

$$-\operatorname{div}(\nabla\phi)\phi \sim \frac{y}{d(d^2 + y^2)}(-d\Delta d), \quad y > 0, x \in \Omega.$$

We are now ready to give the proof of Theorem 4.

**Proof of Theorem 4 parts (i) and (ii).** We assume that  $s \in [\frac{1}{2}, 1)$  or, equivalently,  $a \in (-1, 0]$ . We will use Lemma 3 with the test function  $\phi$  given

$$\phi(x, y) = \begin{cases} (y^2 + d^2)^{-\frac{a}{4}} B\left(\frac{d}{y}\right), & x \in \Omega, y > 0 \\ (y^2 + d^2)^{-\frac{a}{4}} B\left(-\frac{d}{y}\right), & x \in \mathcal{C}\Omega, y > 0. \end{cases}$$

Using Proposition 8 and Lemma 4 we see that all hypotheses of Lemma 3 are satisfied. In particular, we compute

$$\begin{aligned} - \lim_{y \rightarrow 0^+} \left( y^a \frac{\phi_y(x, y)}{\phi(x, y)} \right) &= \frac{1}{d^{1-a}(x)} \lim_{t \rightarrow +\infty} \left( t^{2-a} B'(t) \right) \\ &= \frac{\bar{k}_s}{d^{1-a}(x)}, \quad x \in \Omega. \end{aligned} \tag{3.15}$$

We also have, for  $x \in \Omega$  and  $t = \frac{d}{y} > 0$ ,

$$\begin{aligned} - \operatorname{div}(y^a \nabla\phi) &= -y^a (y^2 + d^2)^{-\frac{a}{4}-1} \left[ (1 + t^2)^2 B'' + (2 - a)t(1 + t^2)B' - \frac{a^2}{4} B \right] \\ &\quad + y^{a+1} (y^2 + d^2)^{-\frac{a}{4}-1} (-\Delta d) \left[ (1 + t^2)B' - \frac{a}{2} tB \right] \\ &= y^{a+1} (y^2 + d^2)^{-\frac{a}{4}-1} (-\Delta d) \left[ (1 + t^2)B' - \frac{a}{2} tB \right], \end{aligned} \tag{3.16}$$



whereas for  $x \in \mathcal{C}\Omega$  and  $t = -\frac{d}{y} < 0$ , we have

$$\begin{aligned} -\operatorname{div}(y^a \nabla \phi) &= -y^a (y^2 + d^2)^{-\frac{a}{4}-1} \left[ (1+t^2)^2 B'' + (2-a)t(1+t^2)B' - \frac{a^2}{4} B \right] \\ &\quad + y^{a+1} (y^2 + d^2)^{-\frac{a}{4}-1} (\Delta d) \left[ (1+t^2)B' - \frac{a}{2} t B \right] \\ &= y^{a+1} (y^2 + d^2)^{-\frac{a}{4}-1} (\Delta d) \left[ (1+t^2)B' - \frac{a}{2} t B \right]. \end{aligned} \quad (3.17)$$

Therefore, under our assumption on  $\Omega$ , it follows from Proposition 8 that

$$-\operatorname{div}(y^a \nabla \phi) \geq 0, \quad x \in \mathbf{R}^n, \quad y > 0.$$

We now use Lemma 3 to get

$$\begin{aligned} \int_0^{+\infty} \int_{\mathbf{R}^n} y^a |\nabla u|^2 \, dx \, dy &\geq \int_0^{+\infty} \int_{\mathbf{R}^n} y^a \left| \nabla u - \frac{\nabla \phi}{\phi} u \right|^2 \, dx \, dy \\ &\quad + \bar{k}_s \int_{\Omega} \frac{u^2(x, 0)}{d^{1-a}(x)} \, dx - \int_0^{+\infty} \int_{\mathbf{R}^n} \frac{\operatorname{div}(y^a \nabla \phi)}{\phi} u^2 \, dx \, dy, \end{aligned} \quad (3.18)$$

from which the trace Hardy inequality follows directly. This relation will also be used later on, in Sections 5 and 6, to obtain the Sobolev term as well.

We next prove the optimality of the Hardy constant. We will work as in Section 2. Let

$$Q[u] := \frac{\int_0^{+\infty} \int_{\mathbf{R}^n} y^a |\nabla u|^2 \, dx \, dy}{\int_{\Omega} \frac{u^2(x, 0)}{d^{1-a}(x)} \, dx} =: \frac{N[u]}{D[u]}. \quad (3.19)$$

We will show that there exists a sequence of functions  $u_\varepsilon$  such that  $\lim_{\varepsilon \rightarrow 0} Q[u_\varepsilon] \leq \bar{k}_s$ , and therefore  $\bar{k}_s$  is the best constant.

We first assume that the boundary of  $\Omega$  is flat in a neighborhood  $U$  of a point  $x_0 \in \partial\Omega$ . The neighborhood of the point  $x_0$  is assumed to contain a ball centered at  $x_0$  with radius, say,  $3\delta$ . Locally around  $x_0$  the boundary is given by  $x_n = 0$ , whereas the interior of  $\Omega$  corresponds to  $x_n > 0$ . We also write  $x = (x', x_n)$ . Clearly, for  $x \in \Omega \cap U$  we have that  $d(x) = x_n$ .

We next define three suitable cutoff functions. Let  $\psi(x') \in C_0^\infty(B_\delta)$ , where  $B_\delta \subset \partial\Omega \subset \mathbf{R}^{n-1}$  is the ball centered at  $x_0$  with radius  $\delta$ . Also, the nonnegative function  $h(x_n) \in C^\infty(\mathbf{R})$  is such that  $h(x_n) = 0$  for  $|x_n| \geq 2\delta$  and  $h(x_n) = 1$  for  $|x_n| \leq \delta$ . We also assume that  $h(x_n)$  is symmetric around  $x_n = 0$ . Finally, let  $\chi(y) \in C_0^\infty(\mathbf{R})$  be such that  $0 \leq \chi(y) \leq 1$ , and  $\chi(y) = 1$  near  $y = 0$ .

We will use the following test function:

$$u_\varepsilon(x', x_n, y) = \chi(y) h(x_n) \psi(x') (y^2 + x_n^2)^{-\frac{a}{4} + \frac{\varepsilon}{4}} B\left(\frac{x_n}{y}\right), \quad x \in \mathbf{R}^n, \quad y > 0. \quad (3.20)$$

Using the asymptotics of  $B(t)$  we easily see that

$$u_\varepsilon(x', x_n, 0) = \begin{cases} h(x_n) \psi(x') x_n^{-\frac{a}{2} + \frac{\varepsilon}{2}}, & x \in \Omega \\ 0, & x \in \mathcal{C}\Omega. \end{cases}$$

We then compute

$$D[u_\varepsilon] = \int_{\mathbf{R}^{n-1}} \psi^2(x') \, dx' \int_0^{+\infty} h^2(x_n) x_n^{-1+\varepsilon} \, dx_n. \quad (3.21)$$

Concerning the numerator, a straightforward calculation shows that

$$\begin{aligned} \left| \nabla \left( (y^2 + x_n^2)^{-\frac{a}{4} + \frac{\varepsilon}{4}} B \left( \frac{x_n}{y} \right) \right) \right|^2 &= \left( -\frac{a}{2} + \frac{\varepsilon}{2} \right)^2 (y^2 + x_n^2)^{-\frac{a}{2} + \frac{\varepsilon}{2} - 1} B^2 \left( \frac{x_n}{y} \right) \\ &\quad + \frac{(x_n^2 + y^2)^{1 - \frac{a}{2} + \frac{\varepsilon}{2}}}{y^4} B'^2 \left( \frac{x_n}{y} \right). \end{aligned}$$

It is then easy to show that

$$\begin{aligned} N[u_\varepsilon] &= O_\varepsilon(1) + \int_{\mathbf{R}^{n-1}} \psi^2(x') \, dx' \int_{\mathbf{R}} \int_0^{+\infty} h^2(x_n) y^a \chi^2(y) \\ &\quad \cdot (y^2 + x_n^2)^{-\frac{a}{2} + \frac{\varepsilon}{2} - 1} \left[ \left( -\frac{a}{2} + \frac{\varepsilon}{2} \right)^2 B^2 \left( \frac{x_n}{y} \right) + \frac{(x_n^2 + y^2)^2}{y^4} B'^2 \left( \frac{x_n}{y} \right) \right] dy \, dx_n. \end{aligned}$$

To estimate the double integral above, we first break the  $x_n$ -integral into two pieces: from minus infinity to zero and from zero to infinity. We then change variables in both pieces by  $t = x_n/y$ , thus going from the  $(x_n, y)$  variables to  $(x_n, t)$ . After elementary calculations, we arrive at

$$\begin{aligned} N[u_\varepsilon] &= O_\varepsilon(1) + \int_{\mathbf{R}^{n-1}} \psi^2(x') \, dx' \int_0^{+\infty} h^2(x_n) x_n^{-1+\varepsilon} \, dx_n \\ &\quad \cdot \int_{-\infty}^{+\infty} \chi^2 \left( \frac{x_n}{|t|} \right) \frac{(1+t^2)^{-1 - \frac{a}{2} + \frac{\varepsilon}{2}}}{|t|^\varepsilon} \left[ (1+t^2)^2 B'^2(t) + \left( -\frac{a}{2} + \frac{\varepsilon}{2} \right)^2 B^2(t) \right] dt. \end{aligned}$$

Forming the quotient, we obtain

$$Q[u_\varepsilon] \leq \int_{-\infty}^{+\infty} \frac{(1+t^2)^{-1 - \frac{a}{2} + \frac{\varepsilon}{2}}}{|t|^\varepsilon} \left[ (1+t^2)^2 B'^2(t) + \frac{(a-\varepsilon)^2}{4} B^2(t) \right] dt + o_\varepsilon(1).$$

We finally send  $\varepsilon$  to zero to get

$$\begin{aligned} \lim_{\varepsilon \rightarrow 0} Q[u_\varepsilon] &\leq \int_{-\infty}^{+\infty} \left[ (1+t^2)^{1 - \frac{a}{2}} B'^2(t) + \frac{a^2}{4} (1+t^2)^{-1 - \frac{a}{2}} B^2(t) \right] dt \\ &= \bar{k}_S; \end{aligned} \quad (3.22)$$

the last equality follows from Proposition 8(iii).

The general case where  $\partial\Omega$  is not flat is treated in the same way as in Section 2.

□

### 4. Some Weighted Hardy Inequalities

In this section we establish some new weighted Hardy inequalities that will play a crucial role in establishing trace Hardy–Sobolev–Maz’ya inequalities.

We first prove the following:

**Lemma 5.** *Let  $\Omega \subset \mathbf{R}^n$  be such that  $-\Delta d(x) \geq 0$  for  $x \in \Omega$ . If  $A, B, \Gamma$  are constants such that  $A + 1 > 0, B + 1 > 0$  and  $2\Gamma < A + B + 2$ , then for all  $v \in C_0^\infty(\mathbf{R}^n \times \mathbf{R})$  there holds*

$$\begin{aligned} & \frac{(B + 1)(B + A + 2 - 2\Gamma^+)}{B + A + 2} \int_0^{+\infty} \int_\Omega \frac{y^A d^B}{(d^2 + y^2)^\Gamma} |v| \, dx \, dy \\ & \leq \int_0^{+\infty} \int_\Omega \frac{y^A d^{B+1}}{(d^2 + y^2)^\Gamma} (-\Delta d) |v| \, dx \, dy + \int_0^{+\infty} \int_\Omega \frac{y^A d^{B+1}}{(d^2 + y^2)^\Gamma} |\nabla v| \, dx \, dy, \end{aligned} \tag{4.1}$$

where  $\Gamma^+ = \max(0, \Gamma)$ .

**Proof.** Integrating by parts in the  $x$ -variables, we compute

$$\begin{aligned} (B + 1) \int_0^{+\infty} \int_\Omega \frac{y^A d^B}{(d^2 + y^2)^\Gamma} |v| \, dx \, dy &= \int_0^{+\infty} \int_\Omega \frac{y^A \nabla d \cdot \nabla d^{B+1}}{(d^2 + y^2)^\Gamma} |v| \, dx \, dy \\ &= \int_0^{+\infty} \int_\Omega \frac{y^A d^{B+1} (-\Delta d)}{(d^2 + y^2)^\Gamma} |v| \, dx \, dy + 2\Gamma \int_0^{+\infty} \int_\Omega \frac{y^A d^{B+2}}{(d^2 + y^2)^{\Gamma+1}} |v| \, dx \, dy \\ &\quad - \int_0^{+\infty} \int_\Omega \frac{y^A d^{B+1}}{(d^2 + y^2)^\Gamma} \nabla d \cdot \nabla_x |v| \, dx \, dy. \end{aligned} \tag{4.2}$$

If  $\Gamma \leq 0$  the result follows easily. In the sequel we consider the case  $\Gamma > 0$ . In the previous calculation there is no boundary term, due to our assumptions. To continue we will estimate the middle term in the right-hand side above. To this end we define the vector field  $\mathbf{F}$  by

$$\mathbf{F}(x, y) := \left( \frac{y^A d^{B+3} \nabla d}{(d^2 + y^2)^{\Gamma+1}}, \frac{y^{A+1} d^{B+2}}{(d^2 + y^2)^{\Gamma+1}} \right). \tag{4.3}$$

We then have

$$\begin{aligned} \int_0^{+\infty} \int_\Omega \operatorname{div} \mathbf{F} |v| \, dx \, dy &= - \int_0^{+\infty} \int_\Omega \mathbf{F} \cdot \nabla |v| \, dx \, dy \\ &\leq \int_0^{+\infty} \int_\Omega |\mathbf{F}| |\nabla v| \, dx \, dy. \end{aligned} \tag{4.4}$$

We note that because of our assumptions  $A + 1 > 0$  and  $B + 1 > 0$ , there are no boundary terms in (4.4). Straightforward calculations show that

$$\operatorname{div} \mathbf{F} = \frac{y^A d^{B+3} (\Delta d)}{(d^2 + y^2)^{\Gamma+1}} + (B + A + 2 - 2\Gamma) \frac{y^A d^{B+2}}{(d^2 + y^2)^{\Gamma+1}}, \tag{4.5}$$

and

$$|\mathbf{F}| = \frac{y^A d^{B+2}}{(d^2 + y^2)^{\Gamma+1/2}} \leq \frac{y^A d^{B+1}}{(d^2 + y^2)^\Gamma}. \quad (4.6)$$

From (4.4)–(4.6) we get

$$\begin{aligned} & (B + A + 2 - 2\Gamma) \int_0^{+\infty} \int_\Omega \frac{y^A d^{B+2}}{(d^2 + y^2)^{\Gamma+1}} |v| \, dx \, dy \\ & \leq \int_0^{+\infty} \int_\Omega \frac{y^A d^{B+3}}{(d^2 + y^2)^{\Gamma+1}} (-\Delta d) |v| \, dx \, dy + \int_0^{+\infty} \int_\Omega \frac{y^A d^{B+1}}{(d^2 + y^2)^\Gamma} |\nabla v| \, dx \, dy. \end{aligned}$$

Combining the above with (4.2) we conclude the proof.  $\square$

We will also need a version of the above lemma in case where  $A + B + 2 = 2\Gamma$ . In this case we have:

**Lemma 6.** *Suppose that  $\Omega \subset \mathbf{R}^n$  is such that  $-\Delta d(x) \geq 0$  for  $x \in \Omega$  and has finite inner radius. If  $A, B$  are constants such that  $A + 1 > 0, B + 1 > 0$ , then for all  $v \in C_0^\infty(\mathbf{R}^n \times \mathbf{R})$  there holds*

$$\begin{aligned} & \frac{B + 1}{A + B + 3} \int_0^{+\infty} \int_\Omega \frac{y^A d^B X^2 |v|}{(d^2 + y^2)^{\frac{A+B+2}{2}}} \, dx \, dy \\ & \leq \int_0^{+\infty} \int_\Omega \frac{y^A d^{B+1} X |v|}{(d^2 + y^2)^{\frac{A+B+2}{2}}} (-\Delta d) \, dx \, dy + \int_0^{+\infty} \int_\Omega \frac{y^A d^{B+1} X |\nabla v|}{(d^2 + y^2)^{\frac{A+B+2}{2}}} \, dx \, dy, \end{aligned} \quad (4.7)$$

where  $X = X\left(\frac{d(x)}{R_{\text{in}}}\right)$  and  $X(t) = (1 - \ln t)^{-1}, 0 < t \leq 1$ .

**Proof.** Integrating by parts in the  $x$ -variables we compute

$$\begin{aligned} & (B + 1) \int_0^{+\infty} \int_\Omega \frac{y^A d^B X^2 |v|}{(d^2 + y^2)^{\frac{A+B+2}{2}}} \, dx \, dy + 2 \int_0^{+\infty} \int_\Omega \frac{y^A d^B X^3 |v|}{(d^2 + y^2)^{\frac{A+B+2}{2}}} \, dx \, dy \\ & \leq \int_0^{+\infty} \int_\Omega \frac{y^A d^{B+1} X^2 (-\Delta d)}{(d^2 + y^2)^{\frac{A+B+2}{2}}} |v| \, dx \, dy \\ & \quad + (A + B + 2) \int_0^{+\infty} \int_\Omega \frac{y^A d^{B+2} X^2}{(d^2 + y^2)^{\frac{A+B+4}{2}}} |v| \, dx \, dy \\ & \quad + \int_0^{+\infty} \int_\Omega \frac{y^A d^{B+1} X^2}{(d^2 + y^2)^{\frac{A+B+2}{2}}} |\nabla v| \, dx \, dy. \end{aligned} \quad (4.8)$$

In the previous calculation there are no boundary terms due to our assumptions. To continue, we will estimate the middle term in the right-hand side above. To this end we define the vector field  $\mathbf{F}$  by

$$\mathbf{F}(x, y) := \left( \frac{y^A d^{B+3} X \nabla d}{(d^2 + y^2)^{\frac{A+B+4}{2}}}, \frac{y^{A+1} d^{B+2} X}{(d^2 + y^2)^{\frac{A+B+4}{2}}} \right). \quad (4.9)$$

We then have

$$\begin{aligned} \int_0^{+\infty} \int_{\Omega} \operatorname{div} \mathbf{F} |v| \, dx \, dy &= - \int_0^{+\infty} \int_{\Omega} \mathbf{F} \cdot \nabla |v| \, dx \, dy \\ &\leq \int_0^{+\infty} \int_{\Omega} |\mathbf{F}| |\nabla v| \, dx \, dy. \end{aligned} \quad (4.10)$$

We note that because of our assumptions  $A + 1 > 0$  and  $B + 1 > 0$ , there are no boundary terms in (4.10). Straightforward calculations show that

$$\operatorname{div} \mathbf{F} = \frac{y^A d^{B+3} X (\Delta d)}{(d^2 + y^2)^{\frac{A+B+4}{2}}} + \frac{y^A d^{B+2} X^2}{(d^2 + y^2)^{\frac{A+B+4}{2}}}, \quad (4.11)$$

and

$$|\mathbf{F}| = \frac{y^A d^{B+2} X}{(d^2 + y^2)^{\frac{A+B+3}{2}}} \leq \frac{y^A d^{B+1} X}{(d^2 + y^2)^{\frac{A+B+2}{2}}}. \quad (4.12)$$

From (4.10)–(4.12) we get

$$\begin{aligned} &\int_0^{+\infty} \int_{\Omega} \frac{y^A d^{B+2} X^2}{(d^2 + y^2)^{\frac{A+B+4}{2}}} |v| \, dx \, dy \\ &\leq \int_0^{+\infty} \int_{\Omega} \frac{y^A d^{B+3} X}{(d^2 + y^2)^{\frac{A+B+4}{2}}} (-\Delta d) |v| \, dx \, dy \\ &\quad + \int_0^{+\infty} \int_{\Omega} \frac{y^A d^{B+1} X}{(d^2 + y^2)^{\frac{A+B+2}{2}}} |\nabla v| \, dx \, dy. \end{aligned}$$

Combining the above with (4.8) we conclude the proof.  $\square$

Without imposing any geometric assumption on  $\Omega$  we have the following result that will also be used later on.

**Lemma 7.** *Let  $\Omega \subset \mathbf{R}^n$ . If  $A, B, \Gamma$  are constants such that  $A + 1 > 0$ ,  $B + 1 > 0$  and  $2\Gamma < A + B + 2$ , then there exist positive constants  $c_1$  and  $c_2$  such that for all  $v \in C_0^\infty(\mathbf{R}^n \times \mathbf{R})$  there holds*

$$\begin{aligned} &\int_0^{+\infty} \int_{\Omega} \frac{y^A d^B}{(d^2 + y^2)^\Gamma} |v| \, dx \, dy \\ &\leq c_1 \int_0^{+\infty} \int_{\Omega} \frac{y^A d^{B+1}}{(d^2 + y^2)^\Gamma} |\nabla v| \, dx \, dy + c_2 \int_0^{+\infty} \int_{\Omega} \frac{y^A d^{B+1}}{(d^2 + y^2)^\Gamma} |v| \, dx \, dy. \end{aligned} \quad (4.13)$$

**Proof.** Here we will use the fact that  $\partial\Omega$  is uniformly Lipschitz. Let  $\{U_i\}$  be a covering of  $\Omega_\varepsilon = \{x \in \Omega : \operatorname{dist}(x, \partial\Omega) < \varepsilon\}$  and let  $\phi_i$  be a partition of unity subordinate to the covering  $\{U_i\}$ . We then have

$$\int_0^{+\infty} \int_{\Omega_\varepsilon} \frac{y^A d^B}{(d^2 + y^2)^\Gamma} |v| \, dx \, dy \leq \sum_{i=1}^{+\infty} \int_0^{+\infty} \int_{\Omega_\varepsilon} \frac{y^A d^B}{(d^2 + y^2)^\Gamma} |\phi_i v| \, dx \, dy.$$

In each  $U_i$  we straighten the boundary and use the equivalence of the distance function to the regularized distance as well as to the difference  $x_n - f_i(x')$  (see [41, section 3.2], or [30, section 12.2]) and obtain

$$\int_0^{+\infty} \int_{\Omega_\varepsilon} \frac{y^A d^B}{(d^2 + y^2)^\Gamma} |\phi_i v| \, dx \, dy \leq C \int_0^{+\infty} \int_{\mathbf{R}_+^n} \frac{y^A t^B}{(t^2 + y^2)^\Gamma} |\tilde{\phi}_i \tilde{v}| \, dx \, dy,$$

for some constant  $C$  independent of  $i$ . We next use Lemma 5 to estimate the right-hand side of this, thus obtaining

$$\begin{aligned} \int_0^{+\infty} \int_{\mathbf{R}_+^n} \frac{y^A t^B}{(t^2 + y^2)^\Gamma} |\tilde{\phi}_i \tilde{v}| \, dx \, dy &\leq c \int_0^{+\infty} \int_{\mathbf{R}_+^n} \frac{y^A t^{B+1}}{(t^2 + y^2)^\Gamma} |\nabla(\tilde{\phi}_i \tilde{v})| \, dx \, dy \\ &\leq c \int_0^{+\infty} \int_{\mathbf{R}_+^n} \frac{y^A t^{B+1}}{(t^2 + y^2)^\Gamma} \tilde{\phi}_i |\nabla \tilde{v}| \, dx \, dy + c \int_0^{+\infty} \int_{\mathbf{R}_+^n} \frac{y^A t^{B+1}}{(t^2 + y^2)^\Gamma} |\nabla \tilde{\phi}_i| |\tilde{v}| \, dx \, dy. \end{aligned}$$

Hence, returning to our original variables, we have

$$\begin{aligned} \int_0^{+\infty} \int_{\Omega_\varepsilon} \frac{y^A d^B}{(d^2 + y^2)^\Gamma} |\phi_i v| \, dx \, dy \\ \leq c \int_0^{+\infty} \int_{\Omega_\varepsilon} \frac{y^A d^{B+1}}{(d^2 + y^2)^\Gamma} \phi_i |\nabla v| \, dx \, dy + c \int_0^{+\infty} \int_{\Omega_\varepsilon} \frac{y^A d^{B+1}}{(d^2 + y^2)^\Gamma} |\nabla \phi_i| |v| \, dx \, dy. \end{aligned}$$

Summing over  $i$  we get

$$\begin{aligned} \int_0^{+\infty} \int_{\Omega_\varepsilon} \frac{y^A d^B}{(d^2 + y^2)^\Gamma} |v| \, dx \, dy \\ \leq C_1 \int_0^{+\infty} \int_{\Omega_\varepsilon} \frac{y^A d^{B+1}}{(d^2 + y^2)^\Gamma} |\nabla v| \, dx \, dy + C_2 \int_0^{+\infty} \int_{\Omega_\varepsilon} \frac{y^A d^{B+1}}{(d^2 + y^2)^\Gamma} |v| \, dx \, dy. \end{aligned}$$

The result then follows easily.  $\square$

When working in the complement of  $\Omega$  we have the following surprising result:

**Lemma 8.** *Let  $\Omega \subset \mathbf{R}^n$ . If  $A, B, \Gamma$  are constants such that  $A + 1 > 0, B + 1 > 0$  and  $2\Gamma < A + B + 2$ , then for all  $v \in C_0^\infty(\mathbf{R}^n \times \mathbf{R})$  there holds*

$$\begin{aligned} (A + 1)(A + B + 2 - 2\Gamma^+) \int_0^{+\infty} \int_{\mathcal{C}\Omega} \frac{y^A d^B}{(d^2 + y^2)^\Gamma} |v| \, dx \, dy \\ \leq 2\Gamma^+ \int_0^{+\infty} \int_{\mathcal{C}\Omega} \frac{y^{A+2} d^{B+1}}{(d^2 + y^2)^{\Gamma+1}} (-\Delta d) |v| \, dx \, dy \\ + (A + B + 2) \int_0^{+\infty} \int_{\mathcal{C}\Omega} \frac{y^{A+1} d^B}{(d^2 + y^2)^\Gamma} |\nabla v| \, dx \, dy, \end{aligned} \tag{4.14}$$

where  $\Gamma^+ = \max(0, \Gamma)$ .

We note that no assumption on the sign of  $-\Delta d$  is required.

**Proof.** Integrating by parts in the  $y$ -variable, we compute

$$(A+1) \int_0^{+\infty} \int_{\mathcal{C}\Omega} \frac{y^A d^B}{(d^2 + y^2)^\Gamma} |v| \, dx \, dy \leq 2\Gamma \int_0^{+\infty} \int_{\mathcal{C}\Omega} \frac{y^{A+2} d^B}{(d^2 + y^2)^{\Gamma+1}} |v| \, dx \, dy + \int_0^{+\infty} \int_{\mathcal{C}\Omega} \frac{y^{A+1} d^B}{(d^2 + y^2)^\Gamma} |\nabla v| \, dx \, dy. \tag{4.15}$$

If  $\Gamma \leq 0$  the result follows easily. In the sequel we consider the case  $\Gamma > 0$ . In the previous calculation there is no boundary term, due to our assumptions. To continue we will estimate the first term in the right-hand side above. To this end we define the vector field  $\mathbf{F}$  by

$$\mathbf{F}(x, y) := \left( \frac{y^{A+2} d^{B+3} \nabla d}{(d^2 + y^2)^{\Gamma+1}}, \frac{y^{A+3} d^B}{(d^2 + y^2)^{\Gamma+1}} \right). \tag{4.16}$$

We then have

$$\begin{aligned} \int_0^{+\infty} \int_{\mathcal{C}\Omega} \operatorname{div} \mathbf{F} |v| \, dx \, dy &= - \int_0^{+\infty} \int_{\mathcal{C}\Omega} \mathbf{F} \cdot \nabla |v| \, dx \, dy \\ &\leq \int_0^{+\infty} \int_{\mathcal{C}\Omega} |\mathbf{F}| |\nabla v| \, dx \, dy. \end{aligned} \tag{4.17}$$

We note that because of our assumptions  $A + 1 > 0$  and  $B + 1 > 0$ , there are no boundary terms in (4.17). Straightforward calculations show that

$$\operatorname{div} \mathbf{F} = \frac{y^{A+2} d^{B+1} (\Delta d)}{(d^2 + y^2)^{\Gamma+1}} + (A + B + 2 - 2\Gamma) \frac{y^{A+2} d^B}{(d^2 + y^2)^{\Gamma+1}}, \tag{4.18}$$

and

$$|\mathbf{F}| = \frac{y^{A+2} d^B}{(d^2 + y^2)^{\Gamma+1/2}} \leq \frac{y^{A+1} d^B}{(d^2 + y^2)^\Gamma}. \tag{4.19}$$

Combining the above we conclude the proof. Again, we note that in all integrations by parts there are no boundary terms, due to our assumptions.  $\square$

As a consequence of Lemma 5 we have:

**Lemma 9.** *Let  $\Omega \subset \mathbf{R}^n$  be such that  $-\Delta d(x) \geq 0$ , for  $x \in \Omega$  and  $w \in C_0^1(\mathbf{R}^n \times \mathbf{R})$ . If  $A, B, \Gamma$  are constants such that  $A + 1 > 0$ ,  $B + 1 > 0$ , and  $2\Gamma < A + B + 2$ , then,*

$$\begin{aligned} &\frac{(B + 1)^2 (B + A + 2 - 2\Gamma^+)^2}{4(B + A + 2)^2} \int_0^{+\infty} \int_{\Omega} \frac{y^A d^B}{(d^2 + y^2)^\Gamma} w^2 \, dx \, dy \\ &\leq \frac{(B + 1)(B + A + 2 - 2\Gamma^+)}{2(B + A + 2)} \int_0^{+\infty} \int_{\Omega} \frac{y^A d^{B+1}}{(d^2 + y^2)^\Gamma} (-\Delta d) w^2 \, dx \, dy \\ &\quad + \int_0^{+\infty} \int_{\Omega} \frac{y^A d^{B+2}}{(d^2 + y^2)^\Gamma} |\nabla w|^2 \, dx \, dy, \end{aligned} \tag{4.20}$$

where  $\Gamma^+ = \max(0, \Gamma)$ .

**Proof.** We apply Lemma 5 to  $v = w^2$ . To conclude, we use Young's inequality in the last term of the right-hand side. We omit the details.  $\square$

In the case where  $A + B + 2 = 2\Gamma$  the  $L^2$  analogue of Lemma 6 reads:

**Lemma 10.** *Suppose that  $\Omega \subset \mathbf{R}^n$  is such that  $-\Delta d(x) \geq 0$  for  $x \in \Omega$  and has finite inner radius. If  $A, B$  are constants such that  $A + 1 > 0, B + 1 > 0$ , then for all  $w \in C_0^\infty(\mathbf{R}^n \times \mathbf{R})$  there holds*

$$\begin{aligned} & \left( \frac{B + 1}{2(A + B + 3)} \right)^2 \int_0^{+\infty} \int_\Omega \frac{y^A d^B X^2}{(d^2 + y^2)^{\frac{A+B+2}{2}}} w^2 \, dx \, dy \\ & \leq \frac{B + 1}{2(A + B + 3)} \int_0^{+\infty} \int_\Omega \frac{y^A d^{B+1} X}{(d^2 + y^2)^{\frac{A+B+2}{2}}} (-\Delta d) w^2 \, dx \, dy \\ & \quad + \int_0^{+\infty} \int_\Omega \frac{y^A d^{B+2}}{(d^2 + y^2)^{\frac{A+B+2}{2}}} |\nabla w|^2 \, dx \, dy, \end{aligned} \tag{4.21}$$

where  $X = X(\frac{d(x)}{R_{\text{in}}})$  and  $X(t) = (1 - \ln t)^{-1}, 0 < t \leq 1$ .

**Proof.** We apply Lemma 6 to  $v = w^2$ . To conclude, we use Young's inequality in the last term of the right-hand side. We omit the details.  $\square$

In the case of half space, a more delicate result is needed. More precisely, we have:

**Lemma 11.** *Let  $v \in C_0^\infty(\mathbf{R}^n \times \mathbf{R})$ . If  $0 < A \leq \frac{1}{2}, B + 1 > 0$ , and  $2\Gamma < A + B + 2$ , then the following inequality holds true:*

$$c_0 \int_0^{+\infty} \int_{\mathbf{R}_+^n} \frac{y^{-A} x_n^B}{(x_n^2 + y^2)^{\Gamma-A}} |v| \, dx \, dy \leq \int_0^{+\infty} \int_{\mathbf{R}_+^n} \frac{y^A x_n^{1+B}}{(x_n^2 + y^2)^\Gamma} |\nabla v| \, dx \, dy, \tag{4.22}$$

where

$$c_0 = \frac{A(B + 1)(B + A + 2 - 2\Gamma^+)}{(A + B + 2)(A + 2B + 2) - 2\Gamma^+(B + 1)}.$$

The same result holds true if we replace  $\mathbf{R}_+^n$  by  $\mathbf{R}_-^n$  with  $|x_n|$  in the place of  $x_n$ .

**Proof.** We will use polar coordinates,  $x_n = r \cos \theta, y = r \sin \theta$ . We first establish the following inequality for the angular derivative.

$$\begin{aligned} A \int_0^{\frac{\pi}{2}} (\sin \theta)^{-A} (\cos \theta)^B |v| \, d\theta & \leq (1 + A + B) \int_0^{\frac{\pi}{2}} (\sin \theta)^{1+A} (\cos \theta)^B |v| \, d\theta \\ & \quad + \int_0^{\frac{\pi}{2}} (\sin \theta)^A (\cos \theta)^{1+B} |v_\theta| \, d\theta. \end{aligned} \tag{4.23}$$

We have

$$\begin{aligned} \frac{d}{d\theta} ((\sin \theta)^A (\cos \theta)^{1+B}) & = A(\sin \theta)^{A-1} (\cos \theta)^{2+B} - (1+B)(\sin \theta)^{A+1} (\cos \theta)^B \\ & = A(\sin \theta)^{A-1} (\cos \theta)^B - (1 + A + B)(\sin \theta)^{A+1} (\cos \theta)^B, \end{aligned}$$



therefore an integration by parts gives:

$$A \int_0^{\frac{\pi}{2}} (\sin \theta)^{A-1} (\cos \theta)^B |v| \, d\theta \leq (1 + A + B) \int_0^{\frac{\pi}{2}} (\sin \theta)^{1+A} (\cos \theta)^B |v| \, d\theta + \int_0^{\frac{\pi}{2}} (\sin \theta)^A (\cos \theta)^{1+B} |v_\theta| \, d\theta.$$

Since  $A \leq \frac{1}{2}$ , we also have that  $(\sin \theta)^{-A} \leq (\sin \theta)^{A-1}$  and (4.23) follows.

We next multiply (4.23) by  $r^{A+B+1-2\Gamma}$  and then integrate over  $(0, \infty)$  to conclude:

$$\begin{aligned} & A \int_0^{+\infty} \int_0^{+\infty} \frac{y^{-A} x_n^B}{(x_n^2 + y^2)^{\Gamma-A}} |v| \, dx_n \, dy \\ & \leq (1 + A + B) \int_0^{+\infty} \int_0^{+\infty} \frac{y^{1+A} x_n^B}{(x_n^2 + y^2)^{\Gamma+\frac{1}{2}}} |v| \, dx_n \, dy \\ & \quad + \int_0^{+\infty} \int_0^{+\infty} \frac{y^A x_n^{1+B}}{(x_n^2 + y^2)^\Gamma} |\nabla v| \, dx_n \, dy \\ & \leq (1 + A + B) \int_0^{+\infty} \int_0^{+\infty} \frac{y^A x_n^B}{(x_n^2 + y^2)^\Gamma} |v| \, dx_n \, dy \\ & \quad + \int_0^{+\infty} \int_0^{+\infty} \frac{y^A x_n^{1+B}}{(x_n^2 + y^2)^\Gamma} |\nabla v| \, dx_n \, dy. \end{aligned} \tag{4.24}$$

We next estimate the first term in the right-hand side by using Lemma 5, that is,

$$\begin{aligned} & \frac{(B + 1)(B + A + 2 - 2\Gamma^+)}{B + A + 2} \int_0^{+\infty} \int_0^{+\infty} \frac{y^A x_n^B}{(x_n^2 + y^2)^\Gamma} |v| \, dx_n \, dy \\ & \leq \int_0^{+\infty} \int_0^{+\infty} \frac{y^A x_n^{B+1}}{(x_n^2 + y^2)^\Gamma} |\nabla v| \, dx_n \, dy. \end{aligned}$$

A further integration in the other variables completes the proof.  $\square$

## 5. Half Space, Trace Hardy and Trace Hardy–Sobolev–Maz’ya Inequalities

Here we will prove the trace Hardy and trace Hardy–Sobolev–Maz’ya inequalities appearing in Theorems 2 and 6. We start with the trace Hardy inequalities.

### 5.1. Half Space, Trace Hardy I and II

In this section we will provide the proof of the trace Hardy inequalities appearing in Theorems 2 and 6.

**Proof of Theorem 2 part (i) and (ii).** The case where  $s \in [\frac{1}{2}, 1)$  is contained in Theorem 1. We next consider the case  $s \in (0, \frac{1}{2})$  or, equivalently,  $a \in (0, 1)$ .

We will use the notation  $x = (x', x_n) \in \mathbf{R}_+^n$  with  $x_n > 0$ . We will use Lemma 1 with the test function  $\phi$  given by

$$\phi(x, y) = x_n^{-\frac{a}{2}} A \left( \frac{y}{x_n} \right), \quad y > 0, \quad x_n > 0, \quad x \in \mathbf{R}_+^n,$$

where  $A$  solves (2.2), (2.3). Using Proposition 7 and Lemma 2 we see that all hypotheses of Lemma 1 are satisfied. In particular, for  $t = \frac{y}{x_n}$  we compute, for  $x \in \mathbf{R}_+^n$ ,

$$- \lim_{y \rightarrow 0^+} \left( y^a \frac{\phi_y(x, y)}{\phi(x, y)} \right) = \frac{\bar{d}_s}{x_n^{1-a}}.$$

We also have

$$-\operatorname{div}(y^a \nabla \phi) = 0, \quad y > 0, \quad x \in \mathbf{R}_+^n.$$

From Lemma 1 we get

$$\begin{aligned} \int_0^{+\infty} \int_{\mathbf{R}_+^n} y^a |\nabla u|^2 \, dx \, dy &\geq \bar{d}_s \int_{\mathbf{R}_+^n} \frac{u^2(x, 0)}{x_n^{1-a}} \, dx \\ &+ \int_0^{+\infty} \int_{\mathbf{R}_+^n} y^a |\nabla u - \frac{\nabla \phi}{\phi} u|^2 \, dx \, dy, \end{aligned} \tag{5.1}$$

from which the trace Hardy inequality follows directly. This relation will be used later on to obtain the Sobolev term, as well.

The optimality of  $\bar{d}_s$  follows by the same test functions given by Equation (2.13) as in the flat case of Theorem 1. The fact that  $a$  covers the full interval  $(-1, 1)$  does not affect the calculations leading to (2.16).  $\square$

**Proof of Theorem 6 part (i).** The case where  $s \in [\frac{1}{2}, 1)$  is contained in Theorem 4. We next consider the case  $s \in (0, \frac{1}{2})$  or, equivalently,  $a \in (0, 1)$ . We will use Lemma 3 with the test function  $\phi$  given

$$\phi(x, y) = (y^2 + x_n^2)^{-\frac{a}{4}} B \left( \frac{x_n}{y} \right), \quad y > 0, \quad x_n \in \mathbf{R}.$$

Using Proposition 8 and Lemma 4 we see that all hypotheses of Lemma 3 are satisfied. In particular, we compute

$$- \lim_{y \rightarrow 0^+} \left( y^a \frac{\phi_y(x, y)}{\phi(x, y)} \right) = \frac{\bar{k}_s}{x_n^{1-a}}, \quad x_n > 0.$$

An easy calculation shows that

$$-\operatorname{div}(y^a \nabla \phi) = 0, \quad x \in \mathbf{R}^n, \quad y > 0.$$

We now use Lemma 3 to get

$$\int_0^{+\infty} \int_{\mathbf{R}^n} y^a |\nabla u|^2 \, dx \, dy \geq \bar{k}_s \int_{\mathbf{R}_+^n} \frac{u^2(x, 0)}{x_n^{1-a}} \, dx + \int_0^{+\infty} \int_{\mathbf{R}^n} y^a \left| \nabla u - \frac{\nabla \phi}{\phi} u \right|^2 \, dx \, dy, \quad (5.2)$$

from which the trace Hardy inequality follows directly. This relation will also be used later on to obtain the Sobolev term, as well.

The optimality of  $\bar{k}_s$  follows by the same test functions given by Equation (3.20) as in the flat case of Theorem 4. The fact that  $a$  covers the full interval  $(-1, 1)$  does not affect the calculations leading to (3.22).  $\square$

### 5.2. Half Space, Trace Hardy–Sobolev–Maz’ya I and II

Here we will give the proof of the trace Hardy–Sobolev–Maz’ya inequalities of Theorems 2 and 6. We will first establish different trace Hardy–Sobolev–Maz’ya inequalities where only the Hardy term appears in the trace, and which are of independent interest.

**Theorem 9.** *Let  $0 < s < 1$  and  $n \geq 2$ . There exists a positive constant  $c$  such that for all  $u \in C_0^\infty(\mathbf{R}_+^\infty \times \mathbf{R})$  there holds*

$$\int_0^{+\infty} \int_{\mathbf{R}_+^n} y^{1-2s} |\nabla_{(x,y)} u(x, y)|^2 \, dx \, dy \geq \bar{d}_s \int_{\mathbf{R}_+^n} \frac{u^2(x, 0)}{x_n^{2s}} \, dx + c \left( \int_0^{+\infty} \int_{\mathbf{R}_+^n} |u(x, y)|^{\frac{2(n+1)}{n-2s}} \, dx \, dy \right)^{\frac{n-2s}{n+1}}. \quad (5.3)$$

with

$$\bar{d}_s := \frac{2\Gamma(1-s)\Gamma^2\left(\frac{3+2s}{4}\right)}{\Gamma^2\left(\frac{3-2s}{4}\right)\Gamma(s)}. \quad (5.4)$$

**Proof of Theorem 9.** From the proof of Theorem 2 we recall the inequality (5.1), that is,

$$\int_0^{+\infty} \int_{\mathbf{R}_+^n} y^a |\nabla u|^2 \, dx \, dy \geq \bar{d}_s \int_{\mathbf{R}_+^n} \frac{u^2(x, 0)}{x_n^{1-a}} \, dx + \int_0^{+\infty} \int_{\mathbf{R}_+^n} y^a \left| \nabla u - \frac{\nabla \phi}{\phi} u \right|^2 \, dx \, dy, \quad (5.5)$$

where  $\phi$  is given by

$$\phi(x, y) = x_n^{-\frac{a}{2}} A\left(\frac{y}{x_n}\right), \quad y > 0, x_n > 0, x \in \mathbf{R}_+^n,$$

and  $A$  solves (2.2), (2.3).

The result will follow after establishing the following inequality:

$$\int_0^{+\infty} \int_{\mathbf{R}_+^n} y^a \left| \nabla u - \frac{\nabla \phi}{\phi} u \right|^2 dx dy \geq c \left( \int_0^{+\infty} \int_{\mathbf{R}_+^n} |u|^{\frac{2(n+1)}{n+a-1}} dx dy \right)^{\frac{n+a-1}{(n+1)}}. \tag{5.6}$$

To this end we start with the inequality, see [33, Theorem 1, section 2.1.6]. For any  $u \in C_0^\infty(\mathbf{R}_+^n \times \mathbf{R})$  we have

$$\int_0^{+\infty} \int_{\mathbf{R}_+^n} y^{\frac{a}{2}} |\nabla u|^2 dx dy \geq c \left( \int_0^{+\infty} \int_{\mathbf{R}_+^n} |u(x, y)|^{\frac{2(n+1)}{2n+a}} dx dy \right)^{\frac{2n+a}{2(n+1)}},$$

with the choice  $u = \phi^{\frac{2n+a}{n+a-1}} v$ . Hence we obtain

$$\begin{aligned} & \int_0^{+\infty} \int_{\mathbf{R}_+^n} y^{\frac{a}{2}} \phi^{\frac{2n+a}{n+a-1}} |\nabla v|^2 dx dy + \frac{2n+a}{n+a-1} \int_0^{+\infty} \int_{\mathbf{R}_+^n} y^{\frac{a}{2}} \phi^{\frac{n+1}{n+a-1}} |\nabla \phi| |v| dx dy \\ & \geq c \left( \int_0^{+\infty} \int_{\mathbf{R}_+^n} |\phi^{\frac{2n+a}{n+a-1}} v|^{\frac{2(n+1)}{2n+a}} dx dy \right)^{\frac{2n+a}{2(n+1)}}. \end{aligned} \tag{5.7}$$

Next we will control the second term of the left-hand side by the first term of the left-hand side. To this end we consider two cases. Suppose first that  $s \in [\frac{1}{2}, 1)$ , that is,  $a \in (-1, 0]$ . Using the asymptotics of Lemma 2 we get that

$$y^{\frac{a}{2}} \phi^{\frac{n+1}{n+a-1}} |\nabla \phi| \sim \frac{y^{\frac{a}{2}} x_n^{\frac{n+1}{n+a-1}}}{(x_n^2 + y^2)^{\frac{(2+a)(2n+a)}{4(n+a-1)}}},$$

whereas

$$y^{\frac{a}{2}} \phi^{\frac{2n+a}{n+a-1}} \sim \frac{y^{\frac{a}{2}} x_n^{\frac{2n+a}{n+a-1}}}{(x_n^2 + y^2)^{\frac{(2+a)(2n+a)}{4(n+a-1)}}}. \tag{5.8}$$

The sought for estimate then is a consequence of Lemma 5 with the choice:  $A = \frac{a}{2}$ ,  $B = \frac{n+1}{n+a-1}$  and  $\Gamma = \frac{(2+a)(2n+a)}{4(n+a-1)}$  taking into account that

$$A + B + 2 - 2\Gamma = \frac{(2-a)(n-1)}{2(n+a-1)} > 0.$$

We next consider the case  $a \in (0, 1)$ . Using again the asymptotics of Lemma 2, this time we have that

$$y^{\frac{a}{2}} \phi^{\frac{n+1}{n+a-1}} |\nabla \phi| \sim \frac{y^{-\frac{a}{2}} x_n^{\frac{n+1}{n+a-1}}}{(x_n^2 + y^2)^{\frac{(2+a)(n+1)}{4(n+a-1)} + \frac{2-a}{4}}},$$

whereas (5.8) remains the same. The sought for estimate now is a consequence of Lemma 11 with the choice  $A = \frac{a}{2}$ ,  $B = \frac{n+1}{n+a-1}$  and  $\Gamma = \frac{(2+a)(2n+a)}{4(n+a-1)}$  taking into account that

$$A + B + 2 - 2\Gamma = \frac{(2-a)(n-1)}{2(n+a-1)} > 0.$$

Therefore, for any  $a \in (-1, 1)$  we arrive at:

$$\int_0^{+\infty} \int_{\mathbf{R}_+^n} y^{\frac{a}{2}} \phi^{\frac{2n+a}{n+a-1}} |\nabla v| \geq c \left( \int_0^{+\infty} \int_{\mathbf{R}_+^n} |\phi^{\frac{2n+a}{n+a-1}} v|^{\frac{2(n+1)}{2n+a}} dx dy \right)^{\frac{2n+a}{2(n+1)}}. \quad (5.9)$$

To continue, we next set in (5.9)  $v = |w|^{\frac{2n+a}{n+a-1}}$  and apply the Schwartz inequality in the left-hand side to conclude, after a simplification,

$$\int_0^{+\infty} \int_{\mathbf{R}_+^n} y^a \phi^2 |\nabla w|^2 dx dy \geq c \left( \int_0^{+\infty} \int_{\mathbf{R}_+^n} |\phi w|^{\frac{2(n+1)}{n+a-1}} dx dy \right)^{\frac{n+a-1}{n+1}}, \quad (5.10)$$

which is equivalent to (5.6).  $\square$

**Proof of Theorem 2 part (iii).** Our starting point now is the following weighted trace Sobolev inequality, see [33, Theorem 1, section 2.1.6],

$$\int_0^{+\infty} \int_{\mathbf{R}_+^n} y^{\frac{a}{2}} |\nabla u| dx dy \geq c \left( \int_{\mathbf{R}_+^n} |u(x, 0)|^{\frac{2n}{2n+a}} dx \right)^{\frac{2n+a}{2n}}, \quad u \in C_0^\infty(\mathbf{R}_+^n \times \mathbf{R}).$$

Again, we set  $u = \phi^{\frac{2n+a}{n+a-1}} v$  to obtain the analogue of (5.7):

$$\begin{aligned} & \int_0^{+\infty} \int_{\mathbf{R}_+^n} y^{\frac{a}{2}} \phi^{\frac{2n+a}{n+a-1}} |\nabla v| dx dy + \frac{2n+a}{n+a-1} \int_0^{+\infty} \int_{\mathbf{R}_+^n} y^{\frac{a}{2}} \phi^{\frac{n+1}{n+a-1}} |\nabla \phi| |v| dx dy \\ & \geq c \left( \int_{\mathbf{R}_+^n} |\phi^{\frac{2n+a}{n+a-1}}(x, 0) v(x, 0)|^{\frac{2n}{2n+a}} dx \right)^{\frac{2n+a}{2n}}. \end{aligned} \quad (5.11)$$

As in the proof of Theorem 9, we control the second term of the left-hand side by the first term of the left-hand side to arrive at

$$\int_0^{+\infty} \int_{\mathbf{R}_+^n} y^{\frac{a}{2}} \phi^{\frac{2n+a}{n+a-1}} |\nabla v| dx dy \geq c \left( \int_{\mathbf{R}_+^n} |\phi^{\frac{2n+a}{n+a-1}}(x, 0) v(x, 0)|^{\frac{2n}{2n+a}} dx \right)^{\frac{2n+a}{2n}}.$$

Again, we set  $v = |w|^{\frac{2n+a}{n+a-1}}$  and apply Schwartz inequality in the left-hand side to arrive at

$$\begin{aligned} & \left( \int_0^{+\infty} \int_{\mathbf{R}_+^n} y^a \phi^2 |\nabla w|^2 dx dy \right)^{\frac{1}{2}} \left( \int_0^{+\infty} \int_{\mathbf{R}_+^n} |\phi w|^{\frac{2(n+1)}{n+a-1}} dx dy \right)^{\frac{1}{2}} \\ & \geq c \left( \int_{\mathbf{R}_+^n} |(\phi w)(x, 0)|^{\frac{2n}{n+a-1}} dx \right)^{\frac{2n+a}{2n}}. \end{aligned}$$

We next use (5.10) to conclude, after a simplification,

$$\int_0^{+\infty} \int_{\mathbf{R}_+^n} y^a \phi^2 |\nabla w|^2 \, dx \, dy \geq c \left( \int_{\mathbf{R}_+^n} |(\phi w)(x, 0)|^{\frac{2n}{n+a-1}} \, dx \right)^{\frac{n+a-1}{n}},$$

which is equivalent to

$$\int_0^{+\infty} \int_{\mathbf{R}_+^n} y^a \left| \nabla u - \frac{\nabla \phi}{\phi} u \right|^2 \, dx \, dy \geq c \left( \int_{\mathbf{R}_+^n} |u(x, 0)|^{\frac{2n}{n+a-1}} \, dx \right)^{\frac{n+a-1}{n}}.$$

Combining this with inequality (5.1) we conclude the proof.  $\square$

We next present a preliminary result which will play an important role towards establishing the Hardy–Sobolev–Maz’ya II of Theorem 6.

**Theorem 10.** *Let  $0 < s < 1$  and  $n \geq 2$ . There exists a positive constant  $c$ , such that for all  $u \in C_0^\infty(\mathbf{R}^n \times \mathbf{R})$  with  $u(x, 0) = 0$ ,  $x \in \mathbf{R}^n$ , there holds*

$$\begin{aligned} \int_0^{+\infty} \int_{\mathbf{R}^n} y^{1-2s} |\nabla_{(x,y)} u(x, y)|^2 \, dx \, dy &\geq \bar{k}_s \int_{\mathbf{R}_+^n} \frac{u^2(x, 0)}{x_n^{2s}} \, dx \\ &+ c \left( \int_0^{+\infty} \int_{\mathbf{R}^n} |u(x, y)|^{\frac{2(n+1)}{n-2s}} \, dx \, dy \right)^{\frac{n-2s}{n+1}}, \end{aligned} \tag{5.12}$$

where

$$\bar{k}_s := \frac{2^{1-2s} \Gamma^2(s + \frac{1}{2}) \Gamma(1 - s)}{\pi \Gamma(s)},$$

is the best constant in (5.12).

**Proof.** From the proof of Theorem 4 we recall the inequality (3.18), that is

$$\begin{aligned} \int_0^{+\infty} \int_{\mathbf{R}^n} y^a |\nabla u|^2 \, dx \, dy &\geq \bar{k}_s \int_{\mathbf{R}_+^n} \frac{u^2(x, 0)}{x_n^{1-a}} \, dx \\ &+ \int_0^{+\infty} \int_{\mathbf{R}^n} y^a \left| \nabla u - \frac{\nabla \phi}{\phi} u \right|^2 \, dx \, dy, \end{aligned} \tag{5.13}$$

where  $\phi$  is given by

$$\phi(x, y) = (y^2 + x_n^2)^{-\frac{a}{4}} B \left( \frac{x_n}{y} \right), \quad y > 0, \quad x_n \in \mathbf{R},$$

and  $B$  solves (3.3), (3.4).

Again, the result will follow after establishing the following inequality:

$$\int_0^{+\infty} \int_{\mathbf{R}^n} y^a \left| \nabla u - \frac{\nabla \phi}{\phi} u \right|^2 \, dx \, dy \geq c \left( \int_0^{+\infty} \int_{\mathbf{R}^n} |u|^{\frac{2(n+1)}{n+a-1}} \, dx \, dy \right)^{\frac{n+a-1}{n+1}} \tag{5.14}$$

To this end we start with the inequality, see [33, Theorem 1, section 2.1.6], valid for any  $u \in C_0^\infty(\mathbf{R}^n \times \mathbf{R})$ :

$$\int_0^{+\infty} \int_{\mathbf{R}^n} y^{\frac{a}{2}} |\nabla u| \, dx \, dy \geq c \left( \int_0^{+\infty} \int_{\mathbf{R}^n} |u(x, y)|^{\frac{2(n+1)}{2n+a}} \, dx \, dy \right)^{\frac{2n+a}{2(n+1)}},$$

with the choice  $u = \phi^{\frac{2n+a}{n+a-1}} v$ . Hence we obtain

$$\begin{aligned} & \int_0^{+\infty} \int_{\mathbf{R}^n} y^{\frac{a}{2}} \phi^{\frac{2n+a}{n+a-1}} |\nabla v| \, dx \, dy + \frac{2n+a}{n+a-1} \int_0^{+\infty} \int_{\mathbf{R}^n} y^{\frac{a}{2}} \phi^{\frac{n+1}{n+a-1}} |\nabla \phi| |v| \, dx \, dy \\ & \geq c \left( \int_0^{+\infty} \int_{\mathbf{R}^n} |\phi^{\frac{2n+a}{n+a-1}} v|^{\frac{2(n+1)}{2n+a}} \, dx \, dy \right)^{\frac{2n+a}{2(n+1)}}. \end{aligned} \tag{5.15}$$

Next we will control the second term of the left-hand side by the first term of the left-hand side. To this end we consider various cases. Suppose, first, that  $s \in [\frac{1}{2}, 1)$  that is  $a \in (-1, 0]$  and  $x \in \mathbf{R}_+^n$ . Using the asymptotics of Lemma 4 we get that

$$y^{\frac{a}{2}} \phi^{\frac{n+1}{n+a-1}} |\nabla \phi| \sim \frac{y^{\frac{a}{2}}}{(x_n^2 + y^2)^{\frac{a(n+1)}{4(n+a-1)} + \frac{a+2}{4}}},$$

whereas

$$y^{\frac{a}{2}} \phi^{\frac{2n+a}{n+a-1}} \sim \frac{y^{\frac{a}{2}}}{(x_n^2 + y^2)^{\frac{a(2n+a)}{4(n+a-1)}}}. \tag{5.16}$$

We now apply Lemma 5 with the choice  $A = \frac{a}{2}$ ,  $B = 0$  and  $\Gamma = \frac{a(n+1)}{4(n+a-1)} + \frac{a+2}{4}$ , taking into account that

$$A + B + 2 - 2\Gamma = \frac{(2-a)(n-1)}{2(n+a-1)} > 0.$$

Thus we get for some positive constant  $c$  that

$$\int_0^{+\infty} \int_{\mathbf{R}_+^n} y^{\frac{a}{2}} \phi^{\frac{2n+a}{n+a-1}} |\nabla v| \, dx \, dy \geq c \int_0^{+\infty} \int_{\mathbf{R}_+^n} y^{\frac{a}{2}} \phi^{\frac{n+1}{n+a-1}} |\nabla \phi| |v| \, dx \, dy. \tag{5.17}$$

We next consider the case  $a \in (0, 1)$ ,  $x \in \mathbf{R}_+^n$ . In this case

$$y^{\frac{a}{2}} \phi^{\frac{n+1}{n+a-1}} |\nabla \phi| \sim \frac{y^{-\frac{a}{2}}}{(x_n^2 + y^2)^{\frac{a(n+1)}{4(n+a-1)} + \frac{2-a}{4}}},$$

whereas

$$y^{\frac{a}{2}} \phi^{\frac{2n+a}{n+a-1}} \sim \frac{y^{\frac{a}{2}}}{(x_n^2 + y^2)^{\frac{a(2n+a)}{4(n+a-1)}}}. \tag{5.18}$$

We now use Lemma 11 with the choice  $A = \frac{a}{2}$ ,  $B = 0$  and  $\Gamma = \frac{1}{2} + \frac{a(2n+a)}{4(n+a-1)}$  taking into account that  $\frac{x_n}{(x_n^2+y^2)^{\frac{1}{2}}} < 1$  and  $A + B + 2 - 2\Gamma = \frac{(2-a)(n-1)}{2(n+a-1)} > 0$ . We then conclude that (5.17) is valid for all  $a \in (-1, 1)$ .

In a similar manner, for all  $a \in (-1, 1)$  and  $x \in \mathbf{R}^n$ , we get that

$$y^{\frac{a}{2}} \phi^{\frac{n+1}{n+a-1}} |\nabla \phi| \sim \frac{y^{-\frac{a}{2} + \frac{(1-a)(n+1)}{n+a-1}}}{(x_n^2 + y^2)^{\frac{(2-a)(2n+a)}{4(n+a-1)}}},$$

whereas

$$y^{\frac{a}{2}} \phi^{\frac{2n+a}{n+a-1}} \sim \frac{y^{\frac{a}{2} + \frac{(1-a)(2n+a)}{n+a-1}}}{(x_n^2 + y^2)^{\frac{(2-a)(2n+a)}{4(n+a-1)}}}. \tag{5.19}$$

This time we use Lemma 8 with  $A = -\frac{a}{2} + \frac{(1-a)(n+1)}{n+a-1}$ ,  $B = 0$  and  $\Gamma = \frac{(2-a)(2n+a)}{4(n+a-1)}$ , noticing that

$$A + B + 2 - 2\Gamma = \frac{(2-a)(n-1)}{2(n+a-1)} > 0,$$

thus obtaining

$$\int_0^{+\infty} \int_{\mathbf{R}^n_-} y^{\frac{a}{2}} \phi^{\frac{2n+a}{n+a-1}} |\nabla v| \, dx \, dy \geq c \int_0^{+\infty} \int_{\mathbf{R}^n_-} y^{\frac{a}{2}} \phi^{\frac{n+1}{n+a-1}} |\nabla \phi| |v| \, dx \, dy. \tag{5.20}$$

Combining (5.17) and (5.20) we obtain the following  $L^1$  Hardy estimate on the whole  $\mathbf{R}^n$ :

$$\int_0^{+\infty} \int_{\mathbf{R}^n} y^{\frac{a}{2}} \phi^{\frac{2n+a}{n+a-1}} |\nabla v| \, dx \, dy \geq c \int_0^{+\infty} \int_{\mathbf{R}^n} y^{\frac{a}{2}} \phi^{\frac{n+1}{n+a-1}} |\nabla \phi| |v| \, dx \, dy. \tag{5.21}$$

Using this in (5.15) we get that

$$\int_0^{+\infty} \int_{\mathbf{R}^n} y^{\frac{a}{2}} \phi^{\frac{2n+a}{n+a-1}} |\nabla v| \, dx \, dy \geq c \left( \int_0^{+\infty} \int_{\mathbf{R}^n} |\phi^{\frac{2n+a}{n+a-1}} v|^{\frac{2(n+1)}{2n+a}} \, dx \, dy \right)^{\frac{2n+a}{2(n+1)}}. \tag{5.22}$$

To continue, we next set in (5.22)  $v = |w|^{\frac{2n+a}{n+a-1}}$  and apply Schwartz inequality in the left-hand side to conclude, after a simplification,

$$\int_0^{+\infty} \int_{\mathbf{R}^n} y^a \phi^2 |\nabla w|^2 \, dx \, dy \geq c \left( \int_0^{+\infty} \int_{\mathbf{R}^n} |\phi w|^{\frac{2(n+1)}{n+a-1}} \, dx \, dy \right)^{\frac{n+a-1}{n+1}}, \tag{5.23}$$

which is equivalent to (5.14). The result then follows.  $\square$

We are now ready to establish the proof of Theorem 6 part (ii).



**Proof of Theorem 6 part (ii).** Again, we will use inequality (5.13). This time the result will follow once we establish the following inequality:

$$\int_0^{+\infty} \int_{\mathbf{R}^n} y^a \left| \nabla u - \frac{\nabla \phi}{\phi} u \right|^2 dx dy \geq c \left( \int_{\mathbf{R}_+^n} |u(x, 0)|^{\frac{2n}{n+a-1}} dx \right)^{\frac{n+a-1}{n}}, \tag{5.24}$$

with  $\phi$  given by

$$\phi(x, y) = (y^2 + x_n^2)^{-\frac{a}{4}} B \left( \frac{x_n}{y} \right), \quad y > 0, \quad x_n \in \mathbf{R},$$

and  $B$  solves (3.3), (3.4).

Our starting point, again, is the following weighted trace Sobolev inequality, see [33, Theorem 1, section 2.1.6], valid for functions  $u \in C_0^\infty(\mathbf{R}^n \times \mathbf{R})$  with  $u(x, 0) = 0, x \in \mathbf{R}_+^n$ :

$$\int_0^{+\infty} \int_{\mathbf{R}^n} y^{\frac{a}{2}} |\nabla u|^2 dx dy \geq c \left( \int_{\mathbf{R}_+^n} |u(x, 0)|^{\frac{2n}{2n+a}} dx \right)^{\frac{2n+a}{2n}}.$$

We set  $u = \phi^{\frac{2n+a}{n+a-1}} v$  to obtain

$$\begin{aligned} & \int_0^{+\infty} \int_{\mathbf{R}^n} y^{\frac{a}{2}} \phi^{\frac{2n+a}{n+a-1}} |\nabla v|^2 dx dy + \frac{2n+a}{n+a-1} \int_0^{+\infty} \int_{\mathbf{R}^n} y^{\frac{a}{2}} \phi^{\frac{n+1}{n+a-1}} |\nabla \phi| |v| dx dy \\ & \geq c \left( \int_{\mathbf{R}_+^n} |\phi^{\frac{2n+a}{n+a-1}}(x, 0) v(x, 0)|^{\frac{2n}{2n+a}} dx \right)^{\frac{2n+a}{2n}}. \end{aligned} \tag{5.25}$$

Combining this with (5.21) we get that

$$\begin{aligned} & \int_0^{+\infty} \int_{\mathbf{R}^n} y^{\frac{a}{2}} \phi^{\frac{2n+a}{n+a-1}} |\nabla v|^2 dx dy \\ & \geq c \left( \int_{\mathbf{R}_+^n} |\phi^{\frac{2n+a}{n+a-1}}(x, 0) v(x, 0)|^{\frac{2n}{2n+a}} dx \right)^{\frac{2n+a}{2n}}. \end{aligned} \tag{5.26}$$

We set  $v = |w|^{\frac{2n+a}{n+a-1}}$  and apply the Schwartz inequality in the left-hand side to arrive at

$$\begin{aligned} & \left( \int_0^{+\infty} \int_{\mathbf{R}^n} y^a \phi^2 |\nabla w|^2 dx dy \right)^{\frac{1}{2}} \left( \int_0^\infty \int_{\mathbf{R}^n} |\phi w|^{\frac{2(n+1)}{n+a-1}} dx dy \right)^{\frac{1}{2}} \\ & \geq c \left( \int_{\mathbf{R}_+^n} |(\phi w)(x, 0)|^{\frac{2n}{n+a-1}} dx \right)^{\frac{2n+a}{2n}}. \end{aligned}$$

We next use the Sobolev inequality (5.23) to conclude, after a simplification,

$$\int_0^{+\infty} \int_{\mathbf{R}^n} y^a \phi^2 |\nabla w|^2 dx dy \geq c \left( \int_{\mathbf{R}_+^n} |(\phi w)(x, 0)|^{\frac{2n}{n+a-1}} dx \right)^{\frac{n+a-1}{n}},$$

which is equivalent to (5.24) and the result follows.  $\square$

## 6. The General Case: Trace Hardy–Sobolev–Maz’ya I and II

### 6.1. Trace Hardy–Sobolev–Maz’ya I

Here we will give the proof of Theorem 1 part (iii). We first establish the following Hardy–Sobolev–Maz’ya, where only the Hardy term appears in the trace term.

**Theorem 11.** *Let  $\frac{1}{2} < s < 1$ ,  $n \geq 2$  and  $\Omega \subsetneq \mathbf{R}^n$  be a uniformly Lipschitz domain with finite inner radius that, in addition, satisfies*

$$-\Delta d(x) \geq 0, \quad x \in \Omega. \tag{6.1}$$

Then there exists a positive constant  $c$  such that for all  $u \in C_0^\infty(\Omega \times \mathbf{R})$  there holds

$$\begin{aligned} \int_0^{+\infty} \int_\Omega y^{1-2s} |\nabla_{(x,y)} u(x, y)|^2 \, dx \, dy &\geq \bar{d}_s \int_\Omega \frac{u^2(x, 0)}{d^{2s}(x)} \, dx \\ &+ c \left( \int_0^{+\infty} \int_\Omega |u(x, y)|^{\frac{2(n+1)}{n-2s}} \, dx \, dy \right)^{\frac{n-2s}{n+1}}. \end{aligned} \tag{6.2}$$

with

$$\bar{d}_s := \frac{2\Gamma(1-s)\Gamma^2\left(\frac{3+2s}{4}\right)}{\Gamma^2\left(\frac{3-2s}{4}\right)\Gamma(s)}. \tag{6.3}$$

**Proof of Theorem 11.** From the proof of Theorem 1 we recall the inequality (2.11), that is,

$$\begin{aligned} \int_0^{+\infty} \int_\Omega y^a |\nabla u|^2 \, dx \, dy &\geq \bar{d}_s \int_\Omega \frac{u^2(x, 0)}{d^{1-a}(x)} \, dx + \int_0^{+\infty} \int_\Omega y^a |\nabla u \\ &- \frac{\nabla\phi}{\phi} u|^2 \, dx \, dy - \int_0^{+\infty} \int_\Omega \frac{\operatorname{div}(y^a \nabla\phi)}{\phi} u^2 \, dx \, dy, \end{aligned} \tag{6.4}$$

where  $\phi$  is given by

$$\phi(x, y) = d^{-\frac{a}{2}}(x)A\left(\frac{y}{d}\right), \quad y > 0, \quad x \in \Omega, \tag{6.5}$$

and  $A$  solves (2.2), (2.3).

The result will follow after establishing the following inequality:

$$\begin{aligned} \int_0^{+\infty} \int_\Omega y^a \left| \nabla u - \frac{\nabla\phi}{\phi} u \right|^2 \, dx \, dy &- \int_0^{+\infty} \int_\Omega \frac{\operatorname{div}(y^a \nabla\phi)}{\phi} u^2 \, dx \, dy \\ &\geq c \left( \int_0^{+\infty} \int_\Omega |u(x, y)|^{\frac{2(n+1)}{n+a-1}} \, dx \, dy \right)^{\frac{n+a-1}{n+1}}. \end{aligned} \tag{6.6}$$

To this end we start with the inequality, see [33, Theorem 1, section 2.1.6], valid for any  $u \in C_0^\infty(\Omega \times \mathbf{R})$ ,

$$\int_0^{+\infty} \int_\Omega y^{\frac{a}{2}} |\nabla u| \, dx \, dy \geq c \left( \int_0^{+\infty} \int_\Omega |u(x, y)|^{\frac{2(n+1)}{2n+a}} \, dx \, dy \right)^{\frac{2n+a}{2(n+1)}},$$

with the choice  $u = \phi^{\frac{2n+a}{n+a-1}} v$ . Hence we obtain

$$\begin{aligned} & \int_0^{+\infty} \int_\Omega y^{\frac{a}{2}} \phi^{\frac{2n+a}{n+a-1}} |\nabla v| \, dx \, dy + \frac{2n+a}{n+a-1} \int_0^{+\infty} \int_\Omega y^{\frac{a}{2}} \phi^{\frac{n+1}{n+a-1}} |\nabla \phi| |v| \, dx \, dy \\ & \geq c \left( \int_0^{+\infty} \int_\Omega |\phi^{\frac{2n+a}{n+a-1}} v|^{\frac{2(n+1)}{2n+a}} \, dx \, dy \right)^{\frac{2n+a}{2(n+1)}}. \end{aligned} \tag{6.7}$$

Next we will control the second term of the left-hand side using Lemma 7. To this end we recall that for  $a \in (-1, 0)$  we have the following asymptotics from Lemma 2:

$$y^{\frac{a}{2}} \phi^{\frac{n+1}{n+a-1}} |\nabla \phi| \sim \frac{y^{\frac{a}{2}} d^{\frac{n+1}{n+a-1}}}{(d^2 + y^2)^{\frac{(2+a)(2n+a)}{4(n+a-1)}}},$$

whereas

$$y^{\frac{a}{2}} \phi^{\frac{2n+a}{n+a-1}} \sim \frac{y^{\frac{a}{2}} d^{\frac{2n+a}{n+a-1}}}{(d^2 + y^2)^{\frac{(2+a)(2n+a)}{4(n+a-1)}}}. \tag{6.8}$$

We then use Lemma 7 with the choice  $A = \frac{a}{2}$ ,  $B = \frac{n+1}{n+a-1}$  and  $\Gamma = \frac{(2+a)(2n+a)}{4(n+a-1)}$ , taking into account that

$$A + B + 2 - 2\Gamma = \frac{(2-a)(n-1)}{2(n+a-1)} > 0,$$

to obtain the estimate

$$\begin{aligned} & \int_0^{+\infty} \int_\Omega \frac{y^{\frac{a}{2}} d^{\frac{n+1}{n+a-1}}}{(d^2 + y^2)^{\frac{(2+a)(2n+a)}{4(n+a-1)}}} |v| \, dx \, dy \\ & \leq C_1 \int_0^{+\infty} \int_\Omega \frac{y^{\frac{a}{2}} d^{\frac{2n+a}{n+a-1}}}{(d^2 + y^2)^{\frac{(2+a)(2n+a)}{4(n+a-1)}}} |\nabla v| \, dx \, dy \\ & \quad + C_2 \int_0^{+\infty} \int_\Omega \frac{y^{\frac{a}{2}} d^{\frac{2n+a}{n+a-1}}}{(d^2 + y^2)^{\frac{(2+a)(2n+a)}{4(n+a-1)}}} |v| \, dx \, dy. \end{aligned}$$

From this and (6.7) we have that

$$\begin{aligned} & \int_0^{+\infty} \int_\Omega y^{\frac{a}{2}} \phi^{\frac{2n+a}{n+a-1}} |\nabla v| \, dx \, dy + \int_0^{+\infty} \int_\Omega y^{\frac{a}{2}} \phi^{\frac{2n+a}{n+a-1}} |v| \, dx \, dy \\ & \geq c \left( \int_0^{+\infty} \int_\Omega |\phi^{\frac{2n+a}{n+a-1}} v|^{\frac{2(n+1)}{2n+a}} \, dx \, dy \right)^{\frac{2n+a}{2(n+1)}}. \end{aligned}$$

To continue, we next set  $v = |w|^{\frac{2n+a}{n+a-1}}$  and apply the Schwartz inequality in the left-hand side. After a simplification, we arrive at:

$$\begin{aligned} & \int_0^{+\infty} \int_{\Omega} y^a \phi^2 |\nabla w|^2 \, dx \, dy + \int_0^{+\infty} \int_{\Omega} y^a \phi^2 w^2 \, dx \, dy \\ & \geq c \left( \int_0^{+\infty} \int_{\Omega} |\phi w|^{\frac{2(n+1)}{n+a-1}} \right)^{\frac{n+a-1}{n+1}} \end{aligned} \quad (6.9)$$

To conclude the proof of the theorem we need the following estimate:

$$\begin{aligned} c \int_0^{+\infty} \int_{\Omega} y^a \phi^2 w^2 \, dx \, dy & \leq \int_0^{+\infty} \int_{\Omega} y^a \phi^2 |\nabla w|^2 \, dx \, dy \\ & \quad - \int_0^{+\infty} \int_{\Omega} \operatorname{div}(y^a \nabla \phi) \phi w^2 \, dx \, dy. \end{aligned} \quad (6.10)$$

It is here that we will use the fact that the domain  $\Omega$  has a finite inner radius. Using Lemma 10 with  $A = a$ ,  $B = 0$ , we obtain that

$$\begin{aligned} c \int_0^{+\infty} \int_{\Omega} \frac{y^a X^2 \left( \frac{d}{R_{\text{in}}} \right)}{(d^2 + y^2)^{\frac{2+a}{2}}} w^2 \, dx \, dy & \leq \int_0^{+\infty} \int_{\Omega} \frac{y^a d^2}{(d^2 + y^2)^{\frac{2+a}{2}}} |\nabla w|^2 \, dx \, dy \\ & \quad - \int_0^{+\infty} \int_{\Omega} \frac{y^a d (\Delta d) X \left( \frac{d}{R_{\text{in}}} \right)}{(d^2 + y^2)^{\frac{2+a}{2}}} w^2 \, dx \, dy, \end{aligned}$$

which implies

$$\begin{aligned} c \int_0^{+\infty} \int_{\Omega} \frac{y^a d^2}{(d^2 + y^2)^{\frac{2+a}{2}}} w^2 \, dx \, dy & \leq \int_0^{+\infty} \int_{\Omega} \frac{y^a d^2}{(d^2 + y^2)^{\frac{2+a}{2}}} |\nabla w|^2 \, dx \, dy \\ & \quad - \int_0^{+\infty} \int_{\Omega} \frac{y^a d (\Delta d)}{(d^2 + y^2)^{\frac{2+a}{2}}} w^2 \, dx \, dy. \end{aligned}$$

Taking into account the asymptotics of  $\phi$ , this is equivalent to (6.10). We omit further details.  $\square$

We are now ready to prove Theorem 1 part (iii).

**Proof of Theorem 1 part (iii).** Again we will use (6.4). The result will follow once we establish

$$\begin{aligned} & \int_0^{+\infty} \int_{\Omega} y^a \left| \nabla u - \frac{\nabla \phi}{\phi} u \right|^2 \, dx \, dy - \int_0^{+\infty} \int_{\Omega} \frac{\operatorname{div}(y^a \nabla \phi)}{\phi} u^2 \, dx \, dy \\ & \geq c \left( \int_{\Omega} |u(x, 0)|^{\frac{2n}{n+a-1}} \, dx \right)^{\frac{n+a-1}{n}}, \end{aligned} \quad (6.11)$$

where  $\phi$  is as in (6.5). To this end we start with the inequality, see [33, Theorem 1, section 2.1.6],

$$\int_0^{+\infty} \int_{\Omega} y^{\frac{a}{2}} |\nabla u| \, dx \, dy \geq c \left( \int_{\Omega} |u(x, 0)|^{\frac{2n}{2n+a}} \, dx \right)^{\frac{2n+a}{2n}}, \quad u \in C_0^\infty(\Omega \times \mathbf{R}),$$

with the choice  $u = \phi^{\frac{2n+a}{n+a-1}} v$ . Hence we obtain

$$\begin{aligned} & \int_0^{+\infty} \int_{\Omega} y^{\frac{a}{2}} \phi^{\frac{2n+a}{n+a-1}} |\nabla v| \, dx \, dy + \frac{2n+a}{n+a-1} \int_0^{+\infty} \int_{\Omega} y^{\frac{a}{2}} \phi^{\frac{n+1}{n+a-1}} |\nabla \phi| |v| \, dx \, dy \\ & \geq c \left( \int_{\Omega} |\phi^{\frac{2n+a}{n+a-1}} v|^{\frac{2n}{2n+a}} \, dx \right)^{\frac{2n+a}{2n}}. \end{aligned} \tag{6.12}$$

Next we will control the second term of the left-hand side exactly as we did in the proof of Theorem 11, to arrive at

$$\begin{aligned} & \int_0^{+\infty} \int_{\Omega} y^{\frac{a}{2}} \phi^{\frac{2n+a}{n+a-1}} |\nabla v| \, dx \, dy + \int_0^{+\infty} \int_{\Omega} y^{\frac{a}{2}} \phi^{\frac{2n+a}{n+a-1}} |v| \, dx \, dy \\ & \geq c \left( \int_{\Omega} |\phi^{\frac{2n+a}{n+a-1}} v(x, 0)|^{\frac{2n}{2n+a}} \, dx \right)^{\frac{2n+a}{2n}}. \end{aligned}$$

To continue, we next set  $v = |w|^{\frac{2n+a}{n+a-1}}$  and apply the Schwartz inequality in the left-hand side to get, after elementary manipulations, that

$$\begin{aligned} & \left( \int_0^{+\infty} \int_{\Omega} |\phi w|^{\frac{2(n+1)}{n+a-1}} \, dx \, dy \right) \left( \int_0^{+\infty} \int_{\Omega} y^a \phi^2 |\nabla w|^2 \, dx \, dy \right) \\ & \quad \times \left( \int_0^{+\infty} \int_{\Omega} |\phi w|^{\frac{2(n+1)}{n+a-1}} \, dx \, dy \right) \left( \int_0^{+\infty} \int_{\Omega} y^a \phi^2 w^2 \, dx \, dy \right) \\ & \geq c \left( \int_{\Omega} |\phi w(x, 0)|^{\frac{2n}{n+a-1}} \, dx \right)^{\frac{2n+a}{n}}. \end{aligned} \tag{6.13}$$

At this point we use Theorem 11 and inequality (6.10) to conclude the result. We omit further details.  $\square$

### 6.2. Trace Hardy–Sobolev–Maz’ya II

Here we will give the proof of Theorem 4 part (iii). We first establish the following Hardy–Sobolev–Maz’ya, where only the Hardy term appears in the trace term.

**Theorem 12.** *Let  $\frac{1}{2} < s < 1$ ,  $n \geq 2$  and  $\Omega \subset \mathbf{R}^n$  be a uniformly Lipschitz and convex domain with finite inner radius. Then there exists a positive constant  $c$  such that for all  $u \in C_0^\infty(\mathbf{R}^n \times \mathbf{R})$  with  $u(x, 0) = 0$  for  $x \in \mathcal{C}\Omega$ , there holds*

$$\begin{aligned} & \int_0^{+\infty} \int_{\mathbf{R}^n} y^{1-2s} |\nabla_{(x,y)} u(x, y)|^2 \, dx \, dy \geq \bar{k}_s \int_{\Omega} \frac{u^2(x, 0)}{d^{2s}(x)} \, dx \\ & \quad + c \left( \int_0^{+\infty} \int_{\mathbf{R}^n} |u(x, y)|^{\frac{2(n+1)}{n-2s}} \, dx \, dy \right)^{\frac{n-2s}{n+1}}, \end{aligned} \tag{6.14}$$

with

$$\bar{k}_s := \frac{2^{1-2s} \Gamma^2(s + \frac{1}{2}) \Gamma(1-s)}{\pi \Gamma(s)}. \tag{6.15}$$

**Proof of Theorem 12.** From the proof of Theorem 4 we recall the inequality (3.18), that is,

$$\begin{aligned} \int_0^{+\infty} \int_{\mathbf{R}^n} y^a |\nabla u|^2 \, dx \, dy &\geq \bar{k}_s \int_{\Omega} \frac{u^2(x, 0)}{d^{1-a}(x)} \, dx + \int_0^{+\infty} \int_{\mathbf{R}^n} y^a \\ &\times \left| \nabla u - \frac{\nabla \phi}{\phi} u \right|^2 \, dx \, dy - \int_0^{+\infty} \int_{\mathbf{R}^n} \frac{\operatorname{div}(y^a \nabla \phi)}{\phi} u^2 \, dx \, dy, \end{aligned} \quad (6.16)$$

where  $\phi$  is given by

$$\phi(x, y) = \begin{cases} (y^2 + d^2)^{-\frac{a}{4}} B\left(\frac{d}{y}\right), & x \in \Omega, \, y > 0 \\ (y^2 + d^2)^{-\frac{a}{4}} B\left(-\frac{d}{y}\right), & x \in \mathcal{C}\Omega, \, y > 0, \end{cases} \quad (6.17)$$

and  $B$  is the solution of the boundary value problem (3.3) and (3.4). The result will follow after establishing the following inequality:

$$\begin{aligned} \int_0^{+\infty} \int_{\mathbf{R}^n} y^a \left| \nabla u - \frac{\nabla \phi}{\phi} u \right|^2 \, dx \, dy - \int_0^{+\infty} \int_{\mathbf{R}^n} \frac{\operatorname{div}(y^a \nabla \phi)}{\phi} u^2 \, dx \, dy \\ \geq c \left( \int_0^{+\infty} \int_{\mathbf{R}^n} |u(x, y)|^{\frac{2(n+1)}{n+a-1}} \, dx \, dy \right)^{\frac{n+a-1}{n+1}}. \end{aligned} \quad (6.18)$$

To this end we start with the inequality, see [33, Theorem 1, section 2.1.6], valid for any  $u \in C_0^\infty(\mathbf{R}^n \times \mathbf{R})$ ,

$$\int_0^{+\infty} \int_{\mathbf{R}^n} y^{\frac{a}{2}} |\nabla u| \, dx \, dy \geq c \left( \int_0^{+\infty} \int_{\mathbf{R}^n} |u(x, y)|^{\frac{2(n+1)}{2n+a}} \, dx \, dy \right)^{\frac{2n+a}{2(n+1)}},$$

with the choice  $u = \phi^{\frac{2n+a}{n+a-1}} v$ . Hence we obtain

$$\begin{aligned} \int_0^{+\infty} \int_{\mathbf{R}^n} y^{\frac{a}{2}} \phi^{\frac{2n+a}{n+a-1}} |\nabla v| \, dx \, dy + \frac{2n+a}{n+a-1} \int_0^{+\infty} \int_{\mathbf{R}^n} y^{\frac{a}{2}} \phi^{\frac{n+1}{n+a-1}} |\nabla \phi| |v| \, dx \, dy \\ \geq c \left( \int_0^{+\infty} \int_{\mathbf{R}^n} |\phi^{\frac{2n+a}{n+a-1}} v|^{\frac{2(n+1)}{2n+a}} \, dx \, dy \right)^{\frac{2n+a}{2(n+1)}}. \end{aligned} \quad (6.19)$$

Again, we want to control the second term of the left-hand side. This time we split the integral into the integral over  $\Omega$  and the integral over  $\mathcal{C}\Omega$ . Concerning the integral over  $\mathcal{C}\Omega$ , we use the asymptotics of  $\phi$  as given by Lemma 4 for  $a \in (-1, 0)$  to get that

$$y^{\frac{a}{2}} \phi^{\frac{n+1}{n+a-1}} |\nabla \phi| \sim \frac{y^{-\frac{a}{2} + \frac{(1-a)(n+1)}{n+a-1}}}{(d^2 + y^2)^{\frac{(2-a)(2n+a)}{4(n+a-1)}}},$$

whereas

$$y^{\frac{a}{2}} \phi^{\frac{2n+a}{n+a-1}} \sim \frac{y^{\frac{a}{2} + \frac{(1-a)(2n+a)}{n+a-1}}}{(d^2 + y^2)^{\frac{(2-a)(2n+a)}{4(n+a-1)}}}.$$

This time we use Lemma 8 with  $A = -\frac{a}{2} + \frac{(1-a)(n+1)}{n+a-1}$ ,  $B = 0$  and  $\Gamma = \frac{(2-a)(2n+a)}{4(n+a-1)}$ , noticing that

$$A + B + 2 - 2\Gamma = \frac{(2-a)(n-1)}{2(n+a-1)} > 0,$$

thus obtaining

$$\int_0^{+\infty} \int_{\mathcal{C}_\Omega} y^{\frac{a}{2}} \phi^{\frac{2n+a}{n+a-1}} |\nabla v| \, dx \, dy \geq c \int_0^{+\infty} \int_{\mathcal{C}_\Omega} y^{\frac{a}{2}} \phi^{\frac{n+1}{n+a-1}} |\nabla \phi| |v| \, dx \, dy, \tag{6.20}$$

where we also used the convexity of  $\Omega$ .

On the other hand, in  $\Omega$  the asymptotics of  $\phi$  are also given by Lemma 4 as follows:

$$y^{\frac{a}{2}} \phi^{\frac{n+1}{n+a-1}} |\nabla \phi| \sim \frac{y^{\frac{a}{2}}}{(d^2 + y^2)^{\frac{a(2n+a)}{4(n+a-1)} + \frac{1}{2}}},$$

whereas

$$y^{\frac{a}{2}} \phi^{\frac{2n+a}{n+a-1}} \sim \frac{y^{\frac{a}{2}}}{(d^2 + y^2)^{\frac{a(2n+a)}{4(n+a-1)}}}.$$

We next use Lemma 7 with the choice  $A = \frac{a}{2}$ ,  $B = 0$  and  $\Gamma = \frac{a(2n+a)}{4(n+a-1)} + \frac{1}{2}$ , taking into account that

$$A + B + 2 - 2\Gamma = \frac{(2-a)(n-1)}{2(n+a-1)} > 0,$$

to obtain the estimate

$$\begin{aligned} & \int_0^{+\infty} \int_{\Omega} \frac{y^{\frac{a}{2}}}{(d^2 + y^2)^{\frac{a(2n+a)}{4(n+a-1)} + \frac{1}{2}}} |v| \, dx \, dy \\ & \leq C_1 \int_0^{+\infty} \int_{\Omega} \frac{y^{\frac{a}{2}} d}{(d^2 + y^2)^{\frac{a(2n+a)}{4(n+a-1)} + \frac{1}{2}}} |\nabla v| \, dx \, dy \\ & \quad + C_2 \int_0^{+\infty} \int_{\Omega} \frac{y^{\frac{a}{2}} d}{(d^2 + y^2)^{\frac{a(2n+a)}{4(n+a-1)} + \frac{1}{2}}} |v| \, dx \, dy \\ & \leq C_1 \int_0^{+\infty} \int_{\Omega} \frac{y^{\frac{a}{2}}}{(d^2 + y^2)^{\frac{a(2n+a)}{4(n+a-1)}}} |\nabla v| \, dx \, dy \\ & \quad + C_2 \int_0^{+\infty} \int_{\Omega} \frac{y^{\frac{a}{2}} d}{(d^2 + y^2)^{\frac{a(2n+a)}{4(n+a-1)} + \frac{1}{2}}} |v| \, dx \, dy. \end{aligned}$$

Equivalently, this can be written as

$$\begin{aligned}
 C \int_0^{+\infty} \int_{\Omega} y^{\frac{a}{2}} \phi^{\frac{n+1}{n+a-1}} |\nabla \phi| |v| \, dx \, dy &\leq \int_0^{+\infty} \int_{\Omega} y^{\frac{a}{2}} \phi^{\frac{2n+a}{n+a-1}} |\nabla v| \, dx \, dy \\
 + \int_0^{+\infty} \int_{\Omega} y^{\frac{a}{2}} \phi^{\frac{2n+a}{n+a-1}} \frac{d}{(d^2 + y^2)^{\frac{1}{2}}} |v| \, dx \, dy. & \tag{6.21}
 \end{aligned}$$

Using (6.20) and (6.21) in (6.19) we arrive at

$$\begin{aligned}
 &\int_0^{+\infty} \int_{\mathbf{R}^n} y^{\frac{a}{2}} \phi^{\frac{2n+a}{n+a-1}} |\nabla v| \, dx \, dy + \int_0^{+\infty} \int_{\Omega} y^{\frac{a}{2}} \frac{d}{(d^2 + y^2)^{\frac{1}{2}}} \phi^{\frac{2n+a}{n+a-1}} |v| \, dx \, dy \\
 &\geq c \left( \int_0^{+\infty} \int_{\mathbf{R}^n} |\phi^{\frac{2n+a}{n+a-1}} v|^{\frac{2(n+1)}{2n+a}} \, dx \, dy \right)^{\frac{2n+a}{2(n+1)}}. \tag{6.22}
 \end{aligned}$$

To continue, we next set  $v = |w|^{\frac{2n+a}{n+a-1}}$  and apply Schwartz inequality in the left-hand side. After a simplification, we arrive at:

$$\begin{aligned}
 &\int_0^{+\infty} \int_{\mathbf{R}^n} y^a \phi^2 |\nabla w|^2 \, dx \, dy + \int_0^{+\infty} \int_{\Omega} \frac{y^a d^2 \phi^2}{d^2 + y^2} w^2 \, dx \, dy \\
 &\geq c \left( \int_0^{+\infty} \int_{\mathbf{R}^n} |\phi w|^{\frac{2(n+1)}{n+a-1}} \right)^{\frac{n+a-1}{n+1}}. \tag{6.23}
 \end{aligned}$$

To conclude the proof of the theorem, it is enough to obtain the following estimate:

$$\begin{aligned}
 c \int_0^{+\infty} \int_{\Omega} \frac{y^a d^2 \phi^2}{d^2 + y^2} w^2 \, dx \, dy &\leq \int_0^{+\infty} \int_{\Omega} y^a \phi^2 |\nabla w|^2 \, dx \, dy \\
 &\quad - \int_0^{+\infty} \int_{\Omega} \operatorname{div}(y^a \nabla \phi) \phi w^2 \, dx \, dy. \tag{6.24}
 \end{aligned}$$

It is here that we will use the fact that the domain  $\Omega$  has a finite inner radius. Using Lemma 10 with  $A = a$ ,  $B = 0$  we obtain that

$$\begin{aligned}
 c \int_0^{+\infty} \int_{\Omega} \frac{y^a X^2 \left( \frac{d}{R_{\text{in}}} \right)}{(d^2 + y^2)^{\frac{2+a}{2}}} w^2 \, dx \, dy &\leq \int_0^{+\infty} \int_{\Omega} \frac{y^a d^2}{(d^2 + y^2)^{\frac{2+a}{2}}} |\nabla w|^2 \, dx \, dy \\
 &\quad - \int_0^{+\infty} \int_{\Omega} \frac{y^a d(\Delta d) X \left( \frac{d}{R_{\text{in}}} \right)}{(d^2 + y^2)^{\frac{2+a}{2}}} w^2 \, dx \, dy,
 \end{aligned}$$

which implies

$$\begin{aligned}
 c \int_0^{+\infty} \int_{\Omega} \frac{y^a d^2}{(d^2 + y^2)^{\frac{2+a}{2}}} w^2 \, dx \, dy &\leq \int_0^{+\infty} \int_{\Omega} \frac{y^a}{(d^2 + y^2)^{\frac{a}{2}}} |\nabla w|^2 \, dx \, dy \\
 &\quad - \int_0^{+\infty} \int_{\Omega} \frac{y^a (\Delta d)}{(d^2 + y^2)^{\frac{1+a}{2}}} w^2 \, dx \, dy.
 \end{aligned}$$

Taking into account the asymptotics of  $\phi$  this is equivalent to (6.24). We omit further details.  $\square$



We are now ready to prove Theorem 4 part (iii).

**Proof of Theorem 4 part (iii).** Again we will use (6.16). The result then will follow once we establish:

$$\begin{aligned} & \int_0^{+\infty} \int_{\mathbf{R}^n} y^a \left| \nabla u - \frac{\nabla \phi}{\phi} u \right|^2 dx dy - \int_0^{+\infty} \int_{\mathbf{R}^n} \frac{\operatorname{div}(y^a \nabla \phi)}{\phi} u^2 dx dy \\ & \geq c \left( \int_{\Omega} |u(x, 0)|^{\frac{2n}{n+a-1}} dx \right)^{\frac{n+a-1}{n}}, \end{aligned} \tag{6.25}$$

where  $\phi$  is as in (6.17). To this end we start again with the inequality,

$$\int_0^{+\infty} \int_{\mathbf{R}^n} y^{\frac{a}{2}} |\nabla u| dx dy \geq c \left( \int_{\Omega} |u(x, 0)|^{\frac{2n}{n+a}} dx \right)^{\frac{2n+a}{2n}},$$

valid for  $u \in C_0^\infty(\mathbf{R}^n \times \mathbf{R})$  with  $u(x, 0) = 0, x \in \mathcal{C}\Omega$ . We apply this to  $u = \phi^{\frac{2n+a}{n+a-1}} v$ . Hence we obtain

$$\begin{aligned} & \int_0^{+\infty} \int_{\mathbf{R}^n} y^{\frac{a}{2}} \phi^{\frac{2n+a}{n+a-1}} |\nabla v| dx dy + \frac{2n+a}{n+a-1} \int_0^{+\infty} \int_{\mathbf{R}^n} y^{\frac{a}{2}} \phi^{\frac{n+1}{n+a-1}} |\nabla \phi| |v| dx dy \\ & \geq c \left( \int_{\Omega} |\phi^{\frac{2n+a}{n+a-1}} v|^{\frac{2n}{2n+a}} dx \right)^{\frac{2n+a}{2n}}. \end{aligned} \tag{6.26}$$

Next we will control the second term of the left-hand side exactly as we did in the proof of Theorem 12, to arrive at

$$\begin{aligned} & \int_0^{+\infty} \int_{\mathbf{R}^n} y^{\frac{a}{2}} \phi^{\frac{2n+a}{n+a-1}} |\nabla v| dx dy + \int_0^{+\infty} \int_{\Omega} \frac{y^{\frac{a}{2}} d}{(d^2 + y^2)^{\frac{1}{2}}} \phi^{\frac{2n+a}{n+a-1}} |v| dx dy \\ & \geq c \left( \int_{\Omega} |\phi^{\frac{2n+a}{n+a-1}} v(x, 0)|^{\frac{2n}{2n+a}} dx \right)^{\frac{2n+a}{2n}}. \end{aligned}$$

To continue, we next set  $v = |w|^{\frac{2n+a}{n+a-1}}$  and apply the Schwartz inequality in the left-hand side to get, after elementary manipulations, that

$$\begin{aligned} & \left( \int_0^{+\infty} \int_{\Omega} |\phi w|^{\frac{2(n+1)}{n+a-1}} dx dy \right) \left( \int_0^{+\infty} \int_{\Omega} y^a \phi^2 |\nabla w|^2 dx dy \right) \\ & \quad \times \left( \int_0^{+\infty} \int_{\Omega} |\phi w|^{\frac{2(n+1)}{n+a-1}} dx dy \right) \left( \int_0^{+\infty} \int_{\Omega} \frac{y^a d^2}{d^2 + y^2} \phi^2 w^2 dx dy \right) \\ & \geq c \left( \int_{\Omega} |\phi w(x, 0)|^{\frac{2n}{n+a-1}} dx \right)^{\frac{2n+a}{n}}. \end{aligned} \tag{6.27}$$

At this point we use Theorem 12 and inequality (6.24) to conclude the result. We omit further details.  $\square$

### 7. The Fractional Laplacians

In this section we will apply the previous results to establish the proofs of Theorems 3, 5 as well as of part (iii) of Theorem 6.

**Proof of Theorem 3.** Parts (i) and (iii) follow from parts (i) and (iii) of Theorem 1, taking into account the relation between the energy of the extended problem and the corresponding one of the fractional Laplacian, see Section 8.1 and, in particular, relation (8.5).

We next prove part (ii). We will use the optimality of the constant  $\bar{d}_s$  of Theorem 1, that is, for each  $\varepsilon > 0$  there exists a  $u_\varepsilon \in C_0^\infty(\Omega \times \mathbf{R})$  such that

$$\bar{d}_s + \varepsilon \geq \frac{\int_0^{+\infty} \int_\Omega y^{1-2s} |\nabla u_\varepsilon|^2 \, dx \, dy}{\int_\Omega \frac{u_\varepsilon^2(x, 0)}{d^{2s}(x)} \, dx},$$

and let  $f_\varepsilon(x) = u_\varepsilon(x, 0)$ . We will show that for some positive constant  $c$ ,

$$d_s + c\varepsilon \geq \frac{((-\Delta)^s f_\varepsilon, f_\varepsilon)_\Omega}{\int_\Omega \frac{f_\varepsilon^2(x)}{d^{2s}(x)} \, dx}. \tag{7.1}$$

To this end, let  $\hat{u}_\varepsilon$  be the solution to the extended problem

$$\begin{aligned} \operatorname{div}(y^{1-2s} \nabla \hat{u}_\varepsilon(x, y)) &= 0, \quad \text{in } \Omega \times (0, \infty), \\ \hat{u}_\varepsilon(x, y) &= 0, \quad x \in \partial\Omega \times (0, \infty), \\ \hat{u}_\varepsilon(x, 0) &= f_\varepsilon(x). \end{aligned}$$

The solution  $\hat{u}_\varepsilon$  minimizes the energy and therefore

$$\int_0^{+\infty} \int_\Omega y^{1-2s} |\nabla \hat{u}_\varepsilon|^2 \, dx \, dy \leq \int_0^{+\infty} \int_\Omega y^{1-2s} |\nabla u_\varepsilon|^2 \, dx \, dy.$$

On the other hand, using (8.5) we have

$$\int_0^{+\infty} \int_\Omega y^{1-2s} |\nabla \hat{u}_\varepsilon|^2 \, dx \, dy = \frac{2^{1-2s} \Gamma(1-s)}{\Gamma(s)} ((-\Delta)^s f_\varepsilon, f_\varepsilon)_\Omega,$$

and (7.1) follows easily with  $c = \frac{\Gamma(s)}{2^{1-2s} \Gamma(1-s)}$ .  $\square$

We next give the proof of Theorem 5

**Proof of Theorem 5.** Parts (i) and (iii) follow from parts (i) and (iii) of Theorem 4, taking into account the relation between the energy of the extended problem and the corresponding one of the fractional Laplacian, see Section 8.2 and, in particular, relations (8.7)–(8.8).

The proof of part (ii) is quite similar to the proof of part (ii) of Theorem 3, the only difference being that the extension problem is now on the whole  $\mathbf{R}^n$ . We omit the details.  $\square$

Finally, estimate (1.33) of part (iii) of Theorem 6 follows at once from part (ii) of Theorem 6 and (8.7). Concerning estimate (1.34), it follows from (1.33), taking into account that for  $x \in \mathbf{R}_+^n$ ,

$$\int_{\mathbf{R}_+^n} \frac{d\xi}{|x - \xi|^{n+2s}} = \frac{\pi^{\frac{n-1}{2}} \Gamma\left(\frac{1+2s}{2}\right)}{2s \Gamma\left(\frac{n+2s}{2}\right)} \frac{1}{x_n^{2s}},$$

see, for example, [7].

## 8. Appendix

### 8.1. Spectral Fractional Laplacian

Let  $\Omega \subset \mathbf{R}^n$  be a bounded domain and let  $\lambda_i$  and  $\phi_i$  be the Dirichlet eigenvalues and eigenfunctions of the Laplacian, that is,  $-\Delta\phi_i = \lambda_i\phi_i$  in  $\Omega$ , with  $\phi_i = 0$  on  $\partial\Omega$ , normalized so that  $\int_{\Omega} \phi_i^2 dx = 1$ . Then, for  $f(x) = \sum c_i\phi_i(x)$  we define

$$(-\Delta)^s f = \sum_{i=1}^{\infty} c_i \lambda_i^s \phi_i, \quad 0 < s < 1. \tag{8.1}$$

We also have

$$((-\Delta)^s f, f)_{\Omega} = \int_{\Omega} f(-\Delta)^s f dx = \sum_{i=1}^{\infty} c_i^2 \lambda_i^s. \tag{8.2}$$

To the function  $f(x)$  we associate the “extended” function  $u(x, y), x \in \Omega, y > 0$ , given by

$$u(x, y) = \sum_{i=1}^{+\infty} c_i \phi_i(x) T(y\sqrt{\lambda_i}),$$

where  $T(t)$  is the energetic solution of the ODE:

$$(t^{1-2s} T'(t))' - t^{1-2s} T(t) = 0, \quad \text{or} \quad T'' + \frac{1-2s}{t} T' - T = 0, \quad t \geq 0. \tag{8.3}$$

The solution of this can be taken from [1, Section 9.6], and is given by

$$T(t) = \frac{2^{1-s}}{\Gamma(s)} t^s K_s(t), \tag{8.4}$$

where  $K_s(t)$  denotes the modified Bessel function of the second kind. The constant factor is chosen in such a way that  $T(0) = 1$ . As a consequence we also have  $u(x, 0) = f(x)$ .

An easy calculation shows that  $\operatorname{div}(y^{1-2s} \nabla(\phi_i(x) T(y\sqrt{\lambda_i}))) = 0$ , from which it follows that  $\operatorname{div}(y^{1-2s} \nabla u) = 0$ . An integration by parts then shows that

$$\begin{aligned}
 \int_0^{+\infty} \int_{\Omega} y^{1-2s} |\nabla u|^2 \, dx \, dy &= \lim_{\tau \rightarrow +\infty} \tau^{1-2s} \int_{\Omega} u(x, \tau) u_y(x, \tau) \, dx \\
 &\quad - \lim_{\tau \rightarrow 0} \tau^{1-2s} \int_{\Omega} u(x, \tau) u_y(x, \tau) \, dx \\
 &= \left[ \lim_{t \rightarrow +\infty} t^{1-2s} T(t) T'(t) - \lim_{t \rightarrow 0} t^{1-2s} T(t) T'(t) \right] \sum_{i=1}^{\infty} \lambda_i^s c_i^2 \\
 &= \frac{2^{1-2s} \Gamma(1-s)}{\Gamma(s)} ((-\Delta)^s f, f)_{\Omega}, \tag{8.5}
 \end{aligned}$$

where we used (8.2) and the fact that

$$\lim_{t \rightarrow +\infty} t^{1-2s} T(t) T'(t) - \lim_{t \rightarrow 0} t^{1-2s} T(t) T'(t) = \frac{2^{1-2s} \Gamma(1-s)}{\Gamma(s)}. \tag{8.6}$$

To prove the above relation, we show that

$$\lim_{t \rightarrow +\infty} t^{1-2s} T(t) T'(t) = 0, \quad - \lim_{t \rightarrow 0} t^{1-2s} T(t) T'(t) = \frac{2^{1-2s} \Gamma(1-s)}{\Gamma(s)}.$$

These two relations are a direct consequence of (8.4) and the following properties of  $K_s(t)$ :

$$\begin{aligned}
 K_s(t) &\sim \frac{\Gamma(s)}{2^{1-s}} t^{-s}, \quad t \rightarrow 0, \quad K_s(t) \sim \sqrt{\frac{\pi}{2t}} e^{-t}, \quad t \rightarrow +\infty, \\
 \frac{d}{dt} (t^s K_s(t)) &= -t^s K_{s-1}(t), \quad K_s(t) = K_{-s}(t).
 \end{aligned}$$

### 8.2. Dirichlet Fractional Laplacian

Let  $u(x, y)$  be the extended function as defined in (1.7)–(1.8). In this subsection we will show the following two relations connecting the energy of the extended problem and the energy of the Dirichlet fractional Laplacian:

$$\begin{aligned}
 \int_0^{+\infty} \int_{\mathbf{R}^n} y^{1-2s} |\nabla u|^2 \, dx \, dy \\
 = \frac{s \Gamma(\frac{n+2s}{2})}{\pi^{\frac{n}{2}} \Gamma(s)} \int_{\mathbf{R}^n} \int_{\mathbf{R}^n} \frac{|f(x) - f(\xi)|^2}{|x - \xi|^{n+2s}} \, dx \, d\xi \tag{8.7}
 \end{aligned}$$

$$\int_0^{+\infty} \int_{\mathbf{R}^n} y^{1-2s} |\nabla u|^2 \, dx \, dy = \frac{2^{1-2s} \Gamma(1-s)}{\Gamma(s)} ((-\Delta)^s f, f)_{\mathbf{R}^n}. \tag{8.8}$$

We will use the Fourier transform in the  $x$ -variables:

$$\hat{u}(\eta, y) = (2\pi)^{-\frac{n}{2}} \int_{\mathbf{R}^n} e^{-ix \cdot \eta} u(x, y) \, dx.$$

The equation  $\operatorname{div}(y^{1-2s} \nabla u(x, y)) = 0$  or, equivalently,  $\Delta_x u + u_{yy} + \frac{a}{y} u_y = 0$  with  $u(x, 0) = f(x)$ , reads as follows when taking the Fourier transform

$$-|\eta|^2 \hat{u} + (\hat{u})_{yy} + \frac{1-2s}{y} (\hat{u})_y = 0, \quad \hat{u}(\eta, 0) = \hat{f}(\eta),$$

and it is satisfied by  $\hat{u}(\eta, y) = \hat{f}(\eta)T(|\eta|y)$ , where  $T$  satisfies (8.3) and is given by Equation (8.4).

Concerning the energies, we have:

$$\begin{aligned} \int_0^{+\infty} y^{1-2s} \int_{\mathbf{R}^n} |\nabla u|^2 \, dx \, dy &= \int_0^{+\infty} y^{1-2s} \int_{\mathbf{R}^n} (|\eta|^2 |\hat{u}|^2 + |\hat{u}_y|^2) \, d\eta \, dy \\ &= \int_0^{+\infty} y^{1-2s} \int_{\mathbf{R}^n} \{|\eta|^2 |\hat{f}|^2 [T^2(|\eta|y) + T'^2(|\eta|y)]\} \, d\eta \, dy \\ &= \left( \int_{\mathbf{R}^n} |\eta|^{2s} |\hat{f}|^2 \, d\eta \right) \left( \int_0^{+\infty} t^{1-2s} [T^2(t) + T'^2(t)] \, dt \right), \end{aligned}$$

where  $t = |\eta|y$ . We next compute the last integral. Multiplying Equation (8.3) by  $T$ , integrating by parts and employing (8.6), we get

$$\int_0^{+\infty} t^{1-2s} [T^2(t) + T'^2(t)] \, dt = t^{1-2s} T(t)T'(t) \Big|_0^\infty = \frac{2^{1-2s} \Gamma(1-s)}{\Gamma(s)}. \tag{8.9}$$

We finally recall the following relation (see, for example, [25, Lemma 3.1]):

$$\begin{aligned} \int_{\mathbf{R}^n} |\eta|^{2s} |\hat{f}|^2 \, d\eta &= \frac{c_{n,s}}{2} \int_{\mathbf{R}^n} \int_{\mathbf{R}^n} \frac{|f(x) - f(\xi)|^2}{|x - \xi|^{n+2s}} \, dx \, d\xi \\ &= \frac{s2^{2s-1} \Gamma\left(\frac{n+2s}{2}\right)}{\pi^{\frac{n}{2}} \Gamma(1-s)} \int_{\mathbf{R}^n} \int_{\mathbf{R}^n} \frac{|f(x) - f(\xi)|^2}{|x - \xi|^{n+2s}} \, dx \, d\xi. \end{aligned} \tag{8.10}$$

Putting together the last three relations, we conclude (8.7).

Finally, taking into account (1.25) we easily obtain (8.8).

*Acknowledgments.* Part of this work was done while SF and AT were visiting the University of Cyprus. LM acknowledges the hospitality of University of Crete during her visit there. AT was partially supported by ELKE grant, University of Crete.

### References

1. ABRAMOWITZ, I., STEGUN, I.: Handbook of Mathematical Functions, with Formulas, Graphs, and Mathematical Tables. *NBS Applied Mathematics Series*, Vol. 55. National Bureau of Standards, Washington, DC, 1964
2. ALBERTI, G., BOUCHITTÉ, G., SEPPECHER, P.: Phase transition with the line-tension effect. *Arch. Ration. Mech. Anal.* **144**(1), 1–46 (1998)
3. ALVINO, A., FERONE, A., VOLPICELLI, R.: Sharp Hardy inequalities in the half space with trace remainder term. *Nonlinear Anal.* **75**(14), 5466–5472 (2012)
4. ANCONA, A.: On strong barriers and an inequality of Hardy for domains in  $\mathbf{R}^n$ . *J. Lond. Math. Soc.* **34**(2), 274–290 (1986)
5. ARMITAGE, D.H., KURAN, U.: The convexity and the superharmonicity of the signed distance function. *Proc. Am. Math. Soc.* **93**(4), 598–600 (1985)
6. BARBATUS, G., FILIPPAS, S., TERTIKAS, A.: A unified approach to improved  $L^p$  Hardy inequalities with best constants. *Trans. Am. Math. Soc.* **356**(6), 2169–2196 (2004)
7. BOGDAN, K., BURDZY, K., CHEN, Z.-Q.: Censored stable processes. *Prob. Theory Relat. Fields* **127**, 89–152 (2003)

8. BOGDAN, K., DYDA, B.: The best constant in a fractional Hardy inequality. *Math. Nachrichten* **284**(5–6), 629–638 (2011)
9. BOURGAIN, J., BREZIS, H., MIRONESCU, P.: Limiting embedding theorems for  $W^{s,p}$  when  $s \uparrow 1$  and applications. *J. Anal. Math.* **87**, 77–101 (2002)
10. BREZIS, H., MARCUS, M.: Hardy's inequalities revisited. Dedicated to Ennio De Giorgi. *Ann. Scuola Norm. Sup. Pisa Cl. Sci.* **4** **25**, 217–237 (1997)
11. CABRÉ, X., CINTI, E.: Energy estimates and 1-D symmetry for nonlinear equations involving the half-Laplacian. *Discrete Contin. Dyn. Syst.* **28**(3), 1179–1206 (2010)
12. CABRÉ, X., TAN, J.: Positive solutions of nonlinear problems involving the square root of the Laplacian. *Adv. Math.* **224**(5), 2052–2093 (2010)
13. CAFFARELLI, L., SILVESTRE, L.: An extension problem related to the fractional Laplacian. *Commun. PDE* **32**, 1245–1260 (2007)
14. CHANG, S.-Y.A., GONZALEZ, M.: Fractional Laplacian in conformal geometry. *Adv. Math.* **226**(2), 1410–1432 (2011)
15. CHEN, Z.-Q., KIM, P., SONG, R.: Two-sided heat kernel estimates for censored stable-like processes. *Probab. Theory Relat. Fields* **146**(3-4), 361–399 (2010)
16. CHEN, Z.-Q., KIM, P., SONG, R.: Heat Kernel Estimates for Dirichlet Fractional Laplacian. *J. Eur. Math. Soc.* **12**(5), 1307–1329 (2010)
17. CHEN, Z.-Q., SONG, R.: Hardy inequality for censored stable processes. *Tohoku Math. J.* **55**(2), 439–450 (2003)
18. DAVIES, E.B.: The Hardy constant. *Q. J. Math.* **46**(2), 417–431 (1995)
19. DAVIES, E.B.: A review of Hardy inequalities. *Oper. Theory Adv. Appl.* **110**, 55–67 (1999)
20. DAVILA, J., DUPAIGNE, L., MONTENEGRO, M.: The extremal solution of a boundary reaction problem. *Commun. Pure Appl. Anal.* **7**(4), 795–817 (2008)
21. DYDA, B.: Fractional Hardy–Sobolev–Maz'ya inequality on balls and halfspaces. Preprint arXiv:1004.5146, 2010
22. EVANS, L.C.: *Partial Differential Equations*. American Mathematical Society, Providence, 1998
23. FILIPPAS, S., MAZ'YA, V., TERTIKAS, A.: Critical Hardy–Sobolev inequalities. *J. Math. Pures Appl.* (9) **87**(1), 37–56 (2007)
24. FRANK, R.L., SEIRINGER, R.: Sharp fractional Hardy inequalities in half-spaces. Around the research of Vladimir Maz'ya. *International Mathematical Series*, Vol. 11 (Eds. A. Laptev), 161–167, 2010
25. FRANK, R.L., LIEB, E.H., SEIRINGER, R.: Hardy–Lieb–Thirring inequalities for fractional Schrödinger operators. *J. Am. Math. Soc.* **21**(4), 925–950 (2008)
26. FRANK, R.L., LOSS, M.: Hardy–Sobolev–Maz'ya inequalities for arbitrary domains. *J. Math. Pures Appl.* (9) **97**, 39–54 (2012)
27. GKIKAS, K.: *Hardy and Hardy–Sobolev Inequalities and their Applications*. Ph.D. thesis, 2011
28. GONZALEZ, M.: Gamma convergence of an energy functional related to the fractional Laplacian. *Calc. Var. Partial Differ. Equ.* **36**(2), 173–210 (2009)
29. KINNUNEN, J., KORTE, R.: Characterizations for the Hardy Inequality. Around the research of Vladimir Maz'ya. *International Mathematical Series*, Vol. 11 (Ed. A. Laptev), 239–254, 2010
30. LEONI, G.: *A First Course in Sobolev Spaces*. Grad. Studies in Math. American Mathematical Society, Providence, 2009
31. LEWIS, R.T., LI, J., LI, Y.Y.: A geometric characterization of a sharp Hardy inequality. *J. Funct. Anal.* **262**(7), 3159–3185 (2012)
32. LOSS, M., SLOANE, C.: Hardy inequalities for fractional integrals on general domains. *J. Funct. Anal.* **259**(6), 1369–1379 (2010)
33. MAZ'YA, V.: *Sobolev Spaces with Applications to Elliptic Partial Differential Equations*. Second, revised and augmented edition. Springer, Heidelberg, 2011

34. MAZ'YA, V., SHAPOSHNIKOVA, T.: On the Bourgain, Brezis, and Mironescu theorem concerning limiting embeddings of fractional Sobolev spaces. *J. Funct. Anal.* **195**(2) 230–238 (2002) [Erratum *J. Funct. Anal.* **201**(1), 298–300 (2003)]
35. NGUYEN, H.M.:  $\Gamma$ -convergence, Sobolev norms and BV functions. *Duke Math. J.* **157**(3), 495–533 (2011)
36. PALATUCCI, G., SAVIN, O., VALDINOCI, E.: *Local and Global Minimizers for a Variational Energy Involving a Fractional Norm*. Preprint, arXiv:1104.1725, 2011
37. PSARADAKIS, G.:  $L^1$  Hardy inequalities with weights. *J. Geom. Anal.* (2012, to appear)
38. SAVIN, O., VALDINOCI, E.: *Density Estimates for a Variational Model Driven by the Gagliardo Norm*. Preprint arXiv:1007.2114, 2010
39. SAVIN, O., VALDINOCI, E.:  $\Gamma$ -convergence for nonlocal phase transitions. *Ann. Inst. H. Poincaré Anal. Non Linéaire* **29**(4), 479–500 (2012)
40. SLOANE, C.A.: A fractional Hardy–Sobolev–Maz'ya inequality on the upper halfspace. *Proc. Am. Math. Soc.* **139**(11) 4003–4016 (2011)
41. STEIN, E.M.: *Singular Integrals and Differentiability Properties of Functions*. Princeton University Press, Princeton, 1970
42. TAN, J.: The Brezis Nirenberg type problem involving the square root of the Laplacian. *Calc. Var. Partial Differ. Equ.* **42**(1–2), 21–41 (2011)

Department of Applied Mathematics,  
University of Crete,  
71409 Heraklion, Greece.  
e-mail: filippas@tem.uoc.gr

and

Dipartimento di Scienze di Base ed Applicate per l'Ingegneria,  
University of Rome "La Sapienza", 00185 Rome, Italy.  
e-mail: moschini@dmmm.uniroma1.it

and

Department of Mathematics,  
University of Crete,  
71409 Heraklion, Greece.  
e-mail: tertikas@math.uoc.gr

(Received January 18, 2012 / Accepted October 5, 2012)

Published online November 21, 2012 – © Springer-Verlag Berlin Heidelberg (2012)